

On a question of Boris Mitjagin

Oleg Reinov

Saint Petersburg State University

Nuclear operators

An operator $T : X \rightarrow Y$ is nuclear if it is of the form

$$Tx = \sum_{k=1}^{\infty} \langle x'_k, x \rangle y_k$$

for all $x \in X$, where $(x'_k) \subset X^*$, $(y_k) \subset Y$, $\sum_k \|x'_k\| \|y_k\| < \infty$.
We use the notation $N(X, Y)$

If T is nuclear, then

$$T : X \rightarrow c_0 \rightarrow l_1 \rightarrow Y.$$



A. Grothendieck, Produits tensoriels topologiques et espaces nucléaires, Mem. Amer. Math. Soc., Volume 16, 1955, 196 + 140.

Let A be a compact operator in H . Then A has the norm convergent expansion

$$A = \sum_{n=1}^N \mu_n(A) (f_n, \cdot) h_n,$$

where $(f_n), (h_n)$ are ONS's, $\mu_1(A) \geq \mu_2(A) \geq \dots > 0$

The $\mu_n(A)$ are called the singular values of A . Notation $s_n(A)$ or just s_n .



Simon B., Trace ideals and their applications, London Math. Soc. Lecture Notes 35, Cambridge University Press, 1979.



$$A \in S_p(H) : \sum s_n^p(A) < \infty, \quad p > 0.$$



$$S_p \circ S_q \subset S_r, \quad 1/r = 1/p + 1/q;$$

$$p, q \in (0, \infty)$$

$$N(H) = S_1(H).$$



$$A \in S_p(H) : \sum s_n^p(A) < \infty, \quad p > 0.$$



$$S_p \circ S_q \subset S_r, \quad 1/r = 1/p + 1/q;$$

$$p, q \in (0, \infty)$$

$$N(H) = S_1(H).$$

s -nuclear operators – Applications de puissance s.éme sommable

- An operator $T : X \rightarrow Y$ is s -nuclear ($0 < s \leq 1$) if it is of the form

$$Tx = \sum_{k=1}^{\infty} \langle x'_k, x \rangle y_k$$

for all $x \in X$, where

$(x'_k) \subset X^*$, $(y_k) \subset Y$, $\sum_k \|x'_k\|^s \|y_k\|^s < \infty$. We use the notation $N_s(X, Y)$.



$$N_p(H) = S_p(H), 0 < p \leq 1.$$

-  R. Oloff, p -normierte Operatorenideale, Beiträge Anal. 4, 105-108 (1972).

s -nuclear operators – Applications de puissance s.éme sommable

- An operator $T : X \rightarrow Y$ is s -nuclear ($0 < s \leq 1$) if it is of the form


$$Tx = \sum_{k=1}^{\infty} \langle x'_k, x \rangle y_k$$

for all $x \in X$, where

$(x'_k) \subset X^*$, $(y_k) \subset Y$, $\sum_k \|x'_k\|^s \|y_k\|^s < \infty$. We use the notation $N_s(X, Y)$.



$$N_p(H) = S_p(H), 0 < p \leq 1.$$

-  R. Oloff, p -normierte Operatorenideale, Beiträge Anal. 4, 105-108 (1972).

On products of nuclear operators

A natural question (due to Boris Mitjagin):

- Is it true that a product of two nuclear operators in Banach spaces can be factored through a trace class (i.e., S_1 -) operator in a Hilbert space?

- By using an example from



Carleman T., Über die Fourierkoeffizienten einer stetigen Funktion, A. M., 41 (1918), 377-384.

it can be shown that

- The answer is negative.

On products of nuclear operators

A natural question (due to Boris Mitjagin):

- Is it true that a product of two nuclear operators in Banach spaces can be factored through a trace class (i.e., S_1 -) operator in a Hilbert space?

- By using an example from



Carleman T., Über die Fourierkoeffizienten einer stetigen Funktion, A. M., 41 (1918), 377-384.

it can be shown that

- The answer is negative.

Explanation

f is Carleman's continuous function:

$$\hat{f} \in l_2 \setminus \cup_{p < 2} l_p.$$

$$T : C \xrightarrow{*f} C.$$

T is nuclear.

Consider the product TT . Note that eigenvalues $(\lambda_k(TT)) \in l_1$ and not better.

Suppose, there is an S_1 -operator $U \in S_1(H)$ so that

$$TT : C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Explanation

f is Carleman's continuous function:

$$\hat{f} \in l_2 \setminus \cup_{p < 2} l_p.$$

$$T : C \overset{*f}{\rightarrow} C.$$

T is nuclear.

Consider the product TT . Note that eigenvalues $(\lambda_k(TT)) \in l_1$ and not better.

Suppose, there is an S_1 -operator $U \in S_1(H)$ so that

$$TT : C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Explanation

f is Carleman's continuous function:

$$\hat{f} \in l_2 \setminus \cup_{p < 2} l_p.$$

$$T : C \overset{*f}{\rightarrow} C.$$

T is nuclear.

Consider the product TT . Note that eigenvalues $(\lambda_k(TT)) \in l_1$ and not better.

Suppose, there is an S_1 -operator $U \in S_1(H)$ so that

$$TT : C \overset{A}{\rightarrow} H \overset{U}{\rightarrow} H \overset{B}{\rightarrow} C.$$

Explanation

f is Carleman's continuous function:

$$\hat{f} \in l_2 \setminus \cup_{p < 2} l_p.$$

$$T : C \xrightarrow{*f} C.$$

T is nuclear.

Consider the product TT . Note that eigenvalues $(\lambda_k(TT)) \in l_1$ and not better.

Suppose, there is an S_1 -operator $U \in S_1(H)$ so that

$$TT : C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Explanation (continued)

Consider

$$H \xrightarrow{B} C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Eigenvalues of UAB = eigenvalues of $TT = BUA$ (and, so, in l_1).
BUT:

$$A \in \Pi_2; \text{ so, } AB \in S_2; U \in S_1.$$

Hence,

$$UAB \in S_{2/3}.$$

Contradiction.

- *Remark.* Sharp fact is that if $V \in NN$, then it factors through an operator $U \in S_2$.

Explanation (continued)

Consider

$$H \xrightarrow{B} C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Eigenvalues of UAB = eigenvalues of $TT = BUA$ (and, so, in l_1).
BUT:

$$A \in \Pi_2; \text{ so, } AB \in S_2; U \in S_1.$$

Hence,

$$UAB \in S_{2/3}.$$

Contradiction.

- *Remark.* Sharp fact is that if $V \in NN$, then it factors through an operator $U \in S_2$.

Explanation (continued)

Consider

$$H \xrightarrow{B} C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Eigenvalues of UAB = eigenvalues of $TT = BUA$ (and, so, in l_1).
BUT:

$$A \in \Pi_2; \text{ so, } AB \in S_2; U \in S_1.$$

Hence,

$$UAB \in S_{2/3}.$$

Contradiction.

- *Remark.* Sharp fact is that if $V \in NN$, then it factors through an operator $U \in S_2$.

Explanation (continued)

Consider

$$H \xrightarrow{B} C \xrightarrow{A} H \xrightarrow{U} H \xrightarrow{B} C.$$

Eigenvalues of UAB = eigenvalues of $TT = BUA$ (and, so, in l_1).
BUT:

$$A \in \Pi_2; \text{ so, } AB \in S_2; U \in S_1.$$

Hence,

$$UAB \in S_{2/3}.$$

Contradiction.

- *Remark.* Sharp fact is that if $V \in NN$, then it factors through an operator $U \in S_2$.

General situation

- Let $\alpha, \beta \in (0, 1]$. If $T \in N_\alpha \circ N_\beta$, then it factors through an S_r -operator, where

$$\frac{1}{r} = \frac{1}{\alpha} + \frac{1}{\beta} - \frac{3}{2}.$$

- Particular cases:

$$\alpha = 1, \beta = \frac{2}{3} \implies r = 1;$$

$$\alpha = 1, \beta = 1 \implies r = 2.$$

- Let $\alpha, \beta \in (0, 1]$. If $T \in N_\alpha \circ N_\beta$, then it factors through an S_r -operator, where

$$\frac{1}{r} = \frac{1}{\alpha} + \frac{1}{\beta} - \frac{3}{2}.$$

- Particular cases:

$$\alpha = 1, \beta = \frac{2}{3} \implies r = 1;$$

$$\alpha = 1, \beta = 1 \implies r = 2.$$

To formulate the theorem, we need a definition:

- The spectrum of A is central-symmetric, if together with any eigenvalue $\lambda \neq 0$ it has the eigenvalue $-\lambda$ of the same multiplicity.

It was proved in a paper by M. I. Zelikin



M. I. Zelikin, "A criterion for the symmetry of a spectrum", Dokl. Akad. Nauk 418 (2008), no. 6, 737-740

- **Theorem.** The spectrum of a nuclear operator A acting on a separable Hilbert space is central-symmetric iff $\text{trace } A^{2n-1} = 0$, $n \in \mathbf{N}$.

To formulate the theorem, we need a definition:

- The spectrum of A is central-symmetric, if together with any eigenvalue $\lambda \neq 0$ it has the eigenvalue $-\lambda$ of the same multiplicity.


It was proved in a paper by M. I. Zelikin



M. I. Zelikin, "A criterion for the symmetry of a spectrum", Dokl. Akad. Nauk 418 (2008), no. 6, 737-740

- **Theorem.** The spectrum of a nuclear operator A acting on a separable Hilbert space is central-symmetric iff $\text{trace } A^{2n-1} = 0, n \in \mathbf{N}$.

We can proof:

- **Theorem.** Let Y be a subspace of a quotient (or a quotient of a subspace) of an L_p -space, $1 \leq p \leq \infty$ and $T \in N_s(Y, Y)$ (s -nuclear), where $1/s = 1 + |1/2 - 1/p|$, The spectrum of T is central-symmetric iff $\text{trace } T^{2n-1} = 0, n = 1, 2, \dots$
- *Remark:* In the theorem "trace" is well defined. The result is sharp.
See also
-  Boris S. Mityagin, Criterion for Z_d -symmetry of a Spectrum of a Compact Operator, arXiv: 1504.05242 [math.FA].


We can proof:

- **Theorem.** Let Y be a subspace of a quotient (or a quotient of a subspace) of an L_p -space, $1 \leq p \leq \infty$ and $T \in N_s(Y, Y)$ (s -nuclear), where $1/s = 1 + |1/2 - 1/p|$, The spectrum of T is central-symmetric iff $\text{trace } T^{2n-1} = 0, n = 1, 2, \dots$
- *Remark:* In the theorem "trace" is well defined. The result is sharp.

See also

-  Boris S. Mityagin, Criterion for Z_d -symmetry of a Spectrum of a Compact Operator, arXiv: 1504.05242 [math.FA].

We can proof:

- **Theorem.** Let Y be a subspace of a quotient (or a quotient of a subspace) of an L_p -space, $1 \leq p \leq \infty$ and $T \in N_s(Y, Y)$ (s -nuclear), where $1/s = 1 + |1/2 - 1/p|$, The spectrum of T is central-symmetric iff $\text{trace } T^{2n-1} = 0, n = 1, 2, \dots$
- *Remark:* In the theorem "trace" is well defined. The result is sharp.
See also
-  Boris S. Mityagin, Criterion for Z_d -symmetry of a Spectrum of a Compact Operator, arXiv: 1504.05242 [math.FA].

Thank you for your attention!