

Discrete Hilbert transforms on sparse sequences

Yurii Belov, Tesfa Y. Mengestie and Kristian Seip

ABSTRACT

Weighted discrete Hilbert transforms $(a_n)_n \mapsto \sum_n a_n v_n / (z - \gamma_n)$ from ℓ_v^2 to a weighted L^2 -space are studied, with $\Gamma = (\gamma_n)$ a sequence of distinct points in the complex plane and $v = (v_n)$ a corresponding sequence of positive numbers. In the special case when $|\gamma_n|$ grows at least exponentially, bounded transforms of this kind are described in terms of a simple relative to the Muckenhoupt (A_2) condition. The special case when z is restricted to another sequence Λ is studied in detail; it is shown that a bounded transform satisfying a certain admissibility condition can be split into finitely many surjective transforms, and precise geometric conditions are found for the invertibility of two such weight transforms. These results can be interpreted as statements about systems of reproducing kernels in certain Hilbert spaces of which de Branges spaces and model subspaces of H^2 are prime examples. In particular, a connection to the Feichtinger conjecture is pointed out. Descriptions of Carleson measures and Riesz bases of normalized reproducing kernels for certain ‘small’ de Branges spaces follow from the results of this paper.

1. Introduction and main results

This paper is concerned with the mapping properties of what we call discrete Hilbert transforms in the complex plane. One aspect of this topic was treated in [7], where all unitary discrete Hilbert transforms were described. When we now turn to questions about boundedness, surjectivity, and invertibility, results of the same generality seem at present out of reach. The results to be presented below are complete only when the discrete Hilbert transforms are defined on particularly sparse sequences. We will nevertheless present the problems in the most general setting, as we believe they merit further investigations. We emphasize the connection with topics such as Carleson measures and Riesz bases of normalized reproducing kernels in Hilbert spaces of analytic functions; this will lead us to the most intriguing general question, namely whether or not the Feichtinger conjecture holds true for systems of reproducing kernels in such spaces.

We begin by assuming that we are given an infinite sequence of distinct points $\Gamma = (\gamma_n)$ in \mathbb{C} and a corresponding sequence of positive numbers $v = (v_n)$, both indexed by the positive integers. We define the weighted Hilbert transform as the map

$$(a_n) \mapsto \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n}, \quad (1.1)$$

which is well defined when (a_n) belongs to $\ell_v^2 = \{(a_n) : \|a\|_v^2 = \sum_{n=1}^{\infty} |a_n|^2 v_n < \infty\}$ and z is a point in the set

$$(\Gamma, v)^* = \left\{ z \in \mathbb{C} : \sum_{n=1}^{\infty} \frac{v_n}{|z - \gamma_n|^2} < \infty \right\}.$$

We denote the transformation defined in (1.1) by $H_{(\Gamma, v)}$ and ask if we may describe those nonnegative measures μ on $(\Gamma, v)^*$ such that $H_{(\Gamma, v)}$ is a bounded map from ℓ_v^2 to $L^2((\Gamma, v)^*, \mu)$. This question is another version of the long-standing problem of finding criteria akin to the Muckenhoupt (A_2) condition for boundedness of two-weight Hilbert transforms (cf. the discussion in the last three chapters of [31]).

The centrepiece of this paper is a solution to the boundedness problem when Γ is exponentially or super-exponentially ‘sparse’, that is, when we have

$$\inf_{n \geq 1} \frac{|\gamma_{n+1}|}{|\gamma_n|} > 1. \quad (1.2)$$

In this case, $(\Gamma, v)^*$ is nonempty and in fact equal to $\mathbb{C} \setminus \Gamma$ if and only if

$$\sum_{n=1}^{\infty} \frac{v_n}{1 + |\gamma_n|^2} < \infty; \quad (1.3)$$

we say that v is an *admissible weight sequence* for Γ if (1.3) holds. When we consider the boundedness problem for such sparse sequences Γ , it is quite natural to partition \mathbb{C} in the following way. Set $\Omega_1 = \{z \in \mathbb{C} : |z| < (|\gamma_1| + |\gamma_2|)/2\}$ and then

$$\Omega_n = \left\{ z \in \mathbb{C} : \frac{|\gamma_{n-1}| + |\gamma_n|}{2} \leq |z| < \frac{|\gamma_n| + |\gamma_{n+1}|}{2} \right\} \quad \text{for } n \geq 2.$$

Our solution to the boundedness problem reads as follows.

THEOREM 1.1. *Suppose that the sequence Γ satisfies the sparseness condition (1.2) and that v is an admissible weight sequence for Γ . If μ is a nonnegative measure on \mathbb{C} with $\mu(\Gamma) = 0$, then the map $H_{(\Gamma, v)}$ is bounded from ℓ_v^2 to $L^2(\mathbb{C}, \mu)$ if and only if*

$$\sup_{n \geq 1} \int_{\Omega_n} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} < \infty \quad (1.4)$$

and

$$\sup_{n \geq 1} \left(\sum_{l=1}^n v_l \sum_{m=n+1}^{\infty} \int_{\Omega_m} \frac{d\mu(z)}{|z|^2} + \sum_{m=1}^n \mu(\Omega_m) \sum_{l=n+1}^{\infty} \frac{v_l}{|\gamma_l|^2} \right) < \infty. \quad (1.5)$$

It should be noted that the condition is symmetric in the two measures $\sum_n v_n \delta_{\gamma_n}$ and μ . This is natural since the theorem also gives a necessary and sufficient condition for the adjoint transformation

$$f \mapsto \left(\int_{\mathbb{C}} \frac{f(z) d\mu(z)}{\bar{z} - \bar{\gamma}_n} \right)_n$$

to be bounded from $L^2(\mathbb{C}, \mu)$ to ℓ_v^2 . The condition (1.5) can be understood as a simple relative to the classical Muckenhoupt (A_2) condition.

Besides its simplicity, the main virtue of Theorem 1.1 is its role as a tool in our study of surjectivity and invertibility of discrete Hilbert transforms. We now turn to the latter topic and require thus that μ be a purely atomic measure. In other words, we are interested in the case when there are a sequence of points $\Lambda = (\lambda_j)$ in $(\Gamma, v)^*$ and a corresponding weight sequence $w = (w_j)$ such that the discrete Hilbert transform

$$(a_n)_n \mapsto \left(\sum_{n=1}^{\infty} \frac{a_n v_n}{\lambda_j - \gamma_n} \right)_j \quad (1.6)$$

is bounded from ℓ_v^2 to ℓ_w^2 . To stress the dependence on the pair (Λ, w) , we denote this transformation by $H_{(\Gamma, v); (\Lambda, w)}$. A duality argument (see the next section) shows that if we

want $H_{(\Gamma,v);(\Lambda,w)}$ to be a surjective map as well, then we must have

$$w_j = \left[\max_{\|a\|_v=1} |(H_{(\Gamma,v)}a)(\lambda_j)| \right]^{-2} = \left(\sum_{n=1}^{\infty} \frac{v_n}{|\lambda_j - \gamma_n|^2} \right)^{-1}, \quad (1.7)$$

up to multiplication by a sequence of positive numbers bounded away from 0 and ∞ . When Γ and v are given and Λ is a sequence in $(\Gamma, v)^*$, we say that the sequence given by (1.7) is the *Bessel weight sequence for Λ with respect to (Γ, v)* . The translation into this discrete setting of Theorem 1.2, to be stated in Section 3, is surprisingly subtle: the sequence Λ splits naturally into three subsequences, one that should be viewed as a perturbation of Γ and then two sequences satisfying only certain ‘extreme’ sparseness conditions.

Our next general question is the following. If $w = (w_j)$ is the Bessel weight sequence for Λ with respect to (Γ, v) and $H_{(\Gamma,v);(\Lambda,w)}$ is a bounded transformation, is it possible to split Λ into a finite union of subsequences Λ' such that, with w' denoting the subsequence of w corresponding to Λ' , each of the transformations $H_{(\Gamma,v);(\Lambda',w')}$ is surjective? As will be explained in the next section, this question would have a positive answer should the Feichtinger conjecture hold true. The following result answers this question when (1.2) holds.

THEOREM 1.2. *Suppose that the sequence Γ satisfies the sparseness condition (1.2) and that v is an admissible weight sequence for Γ . If Λ is a sequence in $\mathbb{C} \setminus \Gamma$, w is the Bessel weight sequence for Λ with respect to (Γ, v) , and the transformation $H_{(\Gamma,v);(\Lambda,w)}$ is bounded, then Λ admits a splitting into a finite union of subsequences such that, for each subsequence Λ' and corresponding subsequence w' of w , the transformation $H_{(\Gamma,v);(\Lambda',w')}$ is surjective.*

We proceed now to our third main result, which is a general statement about invertible discrete Hilbert transforms. The observation that leads to this result, is that the inverse transformation, if it exists, can be identified effectively as another discrete Hilbert transform.

To make a precise statement, we introduce the following terminology. We say that a sequence Λ of distinct points in $(\Gamma, v)^*$ is a *uniqueness sequence for $H_{(\Gamma,v)}$* if there is no nonzero vector a in ℓ_v^2 such that $H_{(\Gamma,v)}a$ vanishes on Λ ; we say that Λ is an *exact uniqueness sequence for $H_{(\Gamma,v)}$* if it is a uniqueness sequence for $H_{(\Gamma,v)}$, but fails to be so on the removal of any one of the points in Λ . If Λ is an exact uniqueness sequence for $H_{(\Gamma,v)}$, then we say that a nontrivial function G defined on $(\Gamma, v)^*$ is a *generating function for Λ* if G vanishes on Λ but, for every λ_j in Λ , there is a nonzero vector $a^{(j)}$ in ℓ_v^2 such that $G(z) = (z - \lambda_j)H_{(\Gamma,v)}a^{(j)}(z)$ for every z in $(\Gamma, v)^*$. It is clear that if a generating function exists, then it is unique up to multiplication by a nonzero constant.

We note that if Λ is an exact uniqueness sequence for $H_{(\Gamma,v)}$, then there exists a unique element $e = (e_n)$ in ℓ_v^2 such that $H_{(\Gamma,v);(\Lambda,w)}e = (1, 0, 0, \dots)$. We set $\nu = (\nu_n)$ and $\varpi = (\varpi_j)$, where

$$\nu_n = v_n |\lambda_1 - \gamma_n|^2 |e_n|^2, \quad (1.8)$$

$\varpi_1 = w_1^{-1}$, and

$$\varpi_j = w_j^{-1} |\lambda_j - \lambda_1|^{-2} \left| \sum_{n=1}^{\infty} \frac{e_n v_n}{(\lambda_j - \gamma_n)^2} \right|^{-2}, \quad (1.9)$$

presuming the series appearing in the latter expression converges absolutely. We see that, plainly, we have absolute convergence of this series whenever Λ admits a generating function.

Our next result reads as follows.

THEOREM 1.3. *Suppose that every exact uniqueness sequence for $H_{(\Gamma,v)}$ admits a generating function. Let Λ be a sequence in $(\Gamma, v)^*$ and let w be the Bessel weight sequence for*

Λ with respect to (Γ, v) . If the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is bounded, then $H_{(\Gamma, v); (\Lambda, w)}$ is an invertible transformation if and only if Λ is an exact uniqueness sequence for $H_{(\Gamma, v)}$ and the transformation $H_{(\Lambda, \varpi); (\Gamma, \nu)}$ is bounded.

Note that when we write that ' $H_{(\Lambda, \varpi); (\Gamma, \nu)}$ is bounded', it is implicitly understood that $\Gamma \subset (\Lambda, \varpi)^*$.

We may observe that if $\gamma_n \rightarrow \infty$ when $n \rightarrow \infty$, then the function

$$\Phi(z) = (z - \lambda_1) \sum_{n=1}^{\infty} \frac{e_n v_n}{z - \gamma_n}, \quad (1.10)$$

and its reciprocal $\Psi = 1/\Phi$ are meromorphic functions in \mathbb{C} , and Φ is then the generating function for Λ . We may then rewrite the expressions for ν and ϖ as

$$\nu_n = v_n |\Psi'(\gamma_n)|^{-2} \quad \text{and} \quad \varpi_j = w_j^{-1} |\Phi'(\lambda_j)|^{-2}. \quad (1.11)$$

Combining Theorem 1.3 with Theorem 1.1, we obtain computable and geometric invertibility criteria when Γ is a sparse sequence as defined by (1.2). To illustrate the nature of these criteria, we highlight the following concrete example, where it is tacitly assumed that (1.2) holds and that $w = (w_n)$ is the Bessel weight sequence for Λ with respect to (Γ, v) .

EXAMPLE 1. Set $V_n = \sum_{m=1}^{n-1} v_m$ for $n > 1$. Assume that both $v_n = o(V_n)$ and $V_n \rightarrow \infty$ when $n \rightarrow \infty$, and write $\Gamma = (\gamma_n)$ and $\Lambda = (\lambda_n)$, with both sequences indexed by the positive integers. Moreover, assume that there exists a positive constant C such that

$$\frac{|\gamma_n - \lambda_n|}{|\gamma_n|} \leq C \frac{v_n}{V_n} \quad (1.12)$$

for every positive integer n .

(1) If, in addition, there is a real constant $c < 1/2$ such that

$$\frac{|\gamma_n|}{|\lambda_n|} - 1 \leq c \frac{v_n}{V_n}$$

for all sufficiently large n , then $H_{(\Gamma, v); (\Lambda, w)}$ is an invertible transformation.

(2) If, on the other hand, there is a positive constant $c > 1/2$ such that

$$\frac{|\gamma_n|}{|\lambda_n|} - 1 \geq c \frac{v_n}{V_n}$$

for all sufficiently large n , then $H_{(\Gamma, v); (\Lambda^{(1)}, w^{(1)})}$ is an invertible transformation, where $\Lambda^{(1)} = (\lambda_2, \lambda_3, \dots)$ and $w^{(1)} = (w_2, w_3, \dots)$.

It follows from (1.12) that $H_{(\Gamma, v); (\Lambda, w)}$ is a bounded transformation, while the respective conditions in (1) and (2) imply that the inverse transformations are bounded, subject to the proviso that, when (2) holds, one point be removed from Λ . This rather puzzling example can be seen as an analogue of the Kadets 1/4 theorem for complex exponentials [21] (cf. the discussion in the next section). We note that if we have the precise relation

$$\frac{|\gamma_n|}{|\lambda_n|} - 1 = \frac{1}{2} \frac{v_n}{V_n},$$

then neither $H_{(\Gamma, v); (\Lambda, w)}$ nor $H_{(\Gamma, v); (\Lambda^{(1)}, w^{(1)})}$ is an invertible transformation. Another curious point is that if we replace the condition that $V_n \rightarrow \infty$ by the assumption that $\sup_n V_n < \infty$, then (1.12) automatically implies that $H_{(\Gamma, v); (\Lambda, w)}$ is an invertible transformation.

An interesting feature of our results for sparse sequences is that invertibility implies that Λ is a perturbation of Γ , in a sense to be made precise. As a consequence, we see that there may

exist bounded transformations $H_{(\Gamma,v);(\Lambda,w)}$ such that no infinite subsequence Λ' of Λ is also a subsequence of another sequence Λ'' for which the associated Hilbert transform is invertible.

Before we turn to the proofs of our three main theorems and a detailed discussion of further results as alluded to above, we will place our study in context by explaining in detail how to translate our problems and findings into statements about systems of reproducing kernels. We are particularly interested in such systems in the distinguished case when the underlying Hilbert space is a de Branges spaces or, more generally, a model subspace of H^2 . The study of such systems has a long history, beginning with the work of Paley and Wiener on systems of nonharmonic Fourier series, and is related to a number of interesting applications. More recently, the advent of the Feichtinger conjecture has given additional impetus to the subject; the special case of the conjecture pertaining to discrete Hilbert transforms appears as an interesting setting in which the ramifications of the general Feichtinger conjecture could be explored.

The present investigation originated in questions raised about systems of reproducing kernels. During the course of our work, we have found it both useful and conceptually appealing to transform the subject into a study of the mapping properties of discrete Hilbert transforms. We have learned to appreciate that the essential difficulties thus seem to appear in a more succinct form.

We close this section with a few words on notation. Throughout this paper, the notation $U(z) \lesssim V(z)$ (or equivalently $V(z) \gtrsim U(z)$) means that there is a constant C such that $U(z) \leq CV(z)$ holds for all z in the set in question, which may be a Hilbert space, a set of complex numbers, or a suitable index set. We write $U(z) \simeq V(z)$ if both $U(z) \lesssim V(z)$ and $V(z) \lesssim U(z)$. As above, a sequence $\Gamma = (\gamma_n)$ of distinct complex numbers will frequently be viewed as a subset of \mathbb{C} . Sometimes we shall need to remove a single point, say γ_m , from such a sequence. The sequence thus obtained will then be written as $\Gamma \setminus \{\gamma_m\}$, where $\{\gamma_m\}$ denotes the set consisting of the single point γ_m .

2. Translation into statements about systems of reproducing kernels

2.1. A class of Hilbert spaces

Let \mathcal{H} be a Hilbert space of complex-valued functions defined on some set Ω in \mathbb{C} . We say that a sequence Λ of distinct points in Ω is a *uniqueness sequence* for \mathcal{H} if no nonzero function in \mathcal{H} vanishes on Λ ; we say that Λ is an *exact uniqueness sequence* for \mathcal{H} if it is a *uniqueness sequence* for \mathcal{H} , but fails to be so on the removal of any one of the points in Λ . If Λ is an exact uniqueness sequence for \mathcal{H} , then we say that a nontrivial function G defined on Ω is a *generating function* for Λ if G vanishes on Λ but, for every λ_j in Λ , there is a nonzero function g_j in \mathcal{H} such that $G(z) = (z - \lambda_j)g_j(z)$ for every z in Ω . It is clear that if a generating function exists, it is unique up to multiplication by a nonzero constant.

We assume that \mathcal{H} satisfies the following three axioms.

(A1) \mathcal{H} has a reproducing kernel κ_λ at every point λ in Ω , that is, the point evaluation functional $\kappa_\lambda : f \rightarrow f(\lambda)$ is continuous in \mathcal{H} for every λ in Ω .

(A2) Every exact uniqueness sequence for \mathcal{H} admits a generating function.

(A3) There exists a sequence of distinct points $\Gamma = (\gamma_n)$ in Ω such that the sequence of normalized reproducing kernels $(\kappa_{\gamma_n} / \|\kappa_{\gamma_n}\|_{\mathcal{H}})$ constitutes a Riesz basis for \mathcal{H} . In addition, there is at least one point z in $\Omega \setminus \Gamma$ for which $\kappa_z \neq 0$.

The second axiom (A2) may be viewed as a weak statement about the possibility of dividing out zeros. To see this, we may observe that (A2) holds trivially if \mathcal{H} has the property that whenever $f(\lambda) = 0$ for some f in \mathcal{H} and λ in Ω , we have that $f(z)/(z - \lambda)$ also belongs to \mathcal{H} . On the other hand, (A2) and (A3) lead to a representation of functions in \mathcal{H} (see below), which

shows that if λ is a point in $\Omega \setminus \bar{\Gamma}$ such that $\kappa_\lambda \neq 0$, then $f(z)/(z - \lambda)$ is in \mathcal{H} whenever f is in \mathcal{H} and $f(\lambda) = 0$. In general, however, this division property need not hold at the accumulation points of Γ when we only assume (A2).

The Riesz basis $(\kappa_{\gamma_n}/\|\kappa_{\gamma_n}\|_{\mathcal{H}})$ has a biorthogonal basis, which we call (g_n) . By axiom (A2), we may write $G(z) = c_n(z - \gamma_n)g_n(z)$ for some nonzero constant c_n . We use the suggestive notation $G'(\gamma_n)$ for the value of $G(z)/(z - \gamma_n)$ at γ_n . The sequence g_n is also a Riesz basis for \mathcal{H} , and therefore every vector h in \mathcal{H} can be written as

$$h(z) = \sum_n h(\gamma_n) \frac{G(z)}{G'(\gamma_n)(z - \gamma_n)}, \quad (2.1)$$

where the sum converges with respect to the norm of \mathcal{H} and

$$\|h\|_{\mathcal{H}}^2 \simeq \sum_n \frac{|h(\gamma_n)|^2}{\|\kappa_{\gamma_n}\|_{\mathcal{H}}^2} < \infty;$$

since point evaluation at every point z is a bounded linear functional, (2.1) also converges pointwise in $\Omega \setminus \Gamma$. By the assumption that $h \mapsto (h(\gamma_n)/\|\kappa_{\gamma_n}\|_{\mathcal{H}})$ is a bijective map from \mathcal{H} to ℓ^2 , this means that

$$\sum_n \frac{\|\kappa_{\gamma_n}\|_{\mathcal{H}}^2}{|G'(\gamma_n)|^2 |z - \gamma_n|^2} < \infty \quad (2.2)$$

whenever z is in $\Omega \setminus \Gamma$. By the last part of axiom (A3), there is at least one such z in $\Omega \setminus \Gamma$. Using the notation

$$v_n = \frac{\|\kappa_{\gamma_n}\|_{\mathcal{H}}^2}{|G'(\gamma_n)|^2},$$

we may therefore deduce from (2.2) that

$$\sum_n \frac{v_n}{1 + |\gamma_n|^2} < \infty. \quad (2.3)$$

We may now change our viewpoint: given a sequence of distinct complex numbers $\Gamma = (\gamma_n)$ and a weight sequence $v = (v_n)$ that satisfy the admissibility condition (2.3), we introduce the space $\mathcal{H}(\Gamma, v)$ consisting of all functions

$$f(z) = \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n}$$

for which

$$\|f\|_{\mathcal{H}(\Gamma, v)}^2 = \sum_{n=1}^{\infty} |a_n|^2 v_n < \infty,$$

assuming that the set $(\Gamma, v)^*$ is nonempty. Thus we obtain the value of a function f in $\mathcal{H}(\Gamma, v)$ at a point z in $(\Gamma, v)^*$ by computing a discrete Hilbert transform.

We say that a nonnegative measure μ on $(\Gamma, v)^*$ is a *Carleson measure* for $\mathcal{H}(\Gamma, v)$ if there is a positive constant C such that

$$\int_{(\Gamma, v)^*} |f(z)|^2 d\mu(z) \leq C \|f\|_{\mathcal{H}(\Gamma, v)}^2$$

holds for every f in $\mathcal{H}(\Gamma, v)$. It is now immediate that μ is a Carleson measure for $\mathcal{H}(\Gamma, v)$ if and only if the map $H_{(\Gamma, v)}$ is bounded from ℓ_v^2 to $L^2((\Gamma, v)^*, \mu)$.

2.2. Systems of reproducing kernels in $\mathcal{H}(\Gamma, v)$

We begin again with some generalities. Let \mathcal{H} be a complex Hilbert space and (f_j) be a sequence of unit vectors in \mathcal{H} . We say that (f_j) is a *Bessel sequence* in \mathcal{H} if there is a positive constant

C such that the inequality

$$\sum_j |\langle f, f_j \rangle_{\mathcal{H}}|^2 \leq C \|f\|_{\mathcal{H}}^2$$

holds for every f in \mathcal{H} . The sequence (f_j) is a *Riesz basic sequence* in \mathcal{H} if there are positive constants A and B such that the inequalities

$$A \sum_j |c_j|^2 \leq \left\| \sum_j c_j f_j \right\|_{\mathcal{H}}^2 \leq B \sum_n |c_n|^2 \quad (2.4)$$

hold for every finite sequence of scalars (c_j) . The Feichtinger conjecture claims that every *Bessel sequence of unit vectors* can be expressed as a *finite union of Riesz basic sequences*.

In an intriguing series of papers [11, 12, 14, 15], it has been revealed that the Feichtinger conjecture is equivalent to the long-standing Kadison–Singer conjecture [22]. We refer to the recent paper [13] for a historical account and an interesting reformulation of the Feichtinger conjecture.

Before turning to the special case of normalized reproducing kernels for $\mathcal{H}(\Gamma, v)$, we record the following consequence of the open mapping theorem [29, p. 73].

LEMMA 2.1. *Suppose that T is a bounded linear transformation from a Hilbert space \mathcal{H}_1 to another Hilbert space \mathcal{H}_2 . Then T is surjective if and only if the adjoint transformation T^* is bounded below.*

If we let T be the map $f \mapsto (\langle f, f_j \rangle_{\mathcal{H}})$, then $T^*(c_j) = \sum_j c_j f_j$. Thus it follows from Lemma 2.1 that (f_j) is a Riesz basic sequence if and only if it is a Bessel sequence for which the moment problem $\langle f, f_j \rangle_{\mathcal{H}} = a_j$ has a solution f in \mathcal{H} for every square-summable sequence (a_j) . We may also set $T = H_{(\Gamma, v); (\Lambda, w)}$ and observe that Lemma 2.1 gives the necessary condition

$$w_j \simeq \left(\sum_{n=1}^{\infty} \frac{v_n}{|\lambda_j - \gamma_n|^2} \right)^{-1}$$

for surjectivity of the transformation $H_{(\Gamma, v); (\Lambda, w)}$.

We return now to the space $\mathcal{H}(\Gamma, v)$. We note that the reproducing kernel of $\mathcal{H}(\Gamma, v)$ at a point z in $(\Gamma, v)^*$ is

$$k_z(\zeta) = \sum_{n=1}^{\infty} \frac{v_n}{(\bar{z} - \bar{\gamma}_n)(\zeta - \gamma_n)};$$

this is a direct consequence of the definition of $\mathcal{H}(\Gamma, v)$. Given a sequence $\Lambda = (\lambda_j)$ in $(\Gamma, v)^*$, we associate with it the corresponding sequence of normalized reproducing kernels $(k_{\lambda_j} / \|k_{\lambda_j}\|_{\mathcal{H}(\Gamma, v)})$. We observe that if w is the Bessel weight sequence for Λ with respect to (Γ, v) , then the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is bounded if and only if the system $(k_{\lambda_j} / \|k_{\lambda_j}\|_{\mathcal{H}(\Gamma, v)})$ is a Bessel sequence in $\mathcal{H}(\Gamma, v)$. Moreover, this transformation is both bounded and surjective if and only if the system $(k_{\lambda_j} / \|k_{\lambda_j}\|_{\mathcal{H}(\Gamma, v)})$ is a Riesz basic sequence in $\mathcal{H}(\Gamma, v)$. Thus Theorem 1.2 shows that the Feichtinger conjecture holds true for systems of normalized reproducing kernels in $\mathcal{H}(\Gamma, v)$ whenever Γ is a sequence satisfying the sparseness condition (1.2).

We finally note that if w is the Bessel weight sequence for Λ with respect to (Γ, v) , then the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is invertible if and only if the system $(k_{\lambda_j} / \|k_{\lambda_j}\|_{\mathcal{H}(\Gamma, v)})$ is a Riesz basis for $\mathcal{H}(\Gamma, v)$. If $\mathcal{H}(\Gamma, v)$ is obtained from a space \mathcal{H} satisfying (A1)–(A3), as described in the previous subsection, then Theorem 1.3 applies. In the special case when $\gamma_n \rightarrow \infty$ as $n \rightarrow \infty$, we may write the meromorphic function defined in (1.10) as $\Phi = F/G$, with G again

denoting the generating function for Γ and F an entire function with a simple zero at each point λ_j . Then the expressions appearing in (1.11) can be restated as

$$\nu_n = v_n \left| \frac{F(\gamma_n)}{G'(\gamma_n)} \right|^2 \quad \text{and} \quad \varpi_j = w_j^{-1} \left| \frac{G(\lambda_j)}{F'(\lambda_j)} \right|^2.$$

2.3. First example: de Branges spaces

The prime examples of Hilbert spaces belonging to the general class described in Subsection 2.1 are found among so-called de Branges spaces and model subspaces of H^2 . To begin with, we note that de Branges spaces may be defined in terms of axioms that are very similar to those introduced above. Indeed, a Hilbert space \mathcal{H} of entire functions that contains a nonzero element is called a *de Branges space* if it satisfies the following three axioms.

(H1) \mathcal{H} has a reproducing kernel κ_λ at every point λ in \mathbb{C} , that is, the point evaluation functional $\kappa_\lambda : f \rightarrow f(\lambda)$ is continuous in \mathcal{H} for every λ in \mathbb{C} .

(H2) If f is in \mathcal{H} and $f(\lambda) = 0$ for some point λ in \mathbb{C} , then $f(z)(z - \bar{\lambda})/(z - \lambda)$ is in \mathcal{H} and has the same norm as f .

(H3) The function $\overline{f(\bar{z})}$ belongs to \mathcal{H} whenever f belongs to \mathcal{H} , and it has the same norm as f .

The general reference for de Branges spaces is the book [10]. The leading example of a de Branges space is the Paley–Wiener space, which consists of those entire functions of exponential type at most π that are square summable when restricted to the real line.

A space \mathcal{H} that satisfies (H1)–(H3), will in particular satisfy (A1)–(A3) with $\Omega = \mathbb{C}$. Indeed, we observe that then (H1) and (A1) coincide, and it is also plain that (H2) implies (A2). One of the basic results in de Branges’s theory is that a space that satisfies (H1)–(H3), will have an orthogonal basis consisting of reproducing kernels κ_{γ_n} with $\Gamma = (\gamma_n)$ being a sequence of real points. Thus, in particular, (H1)–(H3) imply that our third general axiom (A3) holds. In the case of the Paley–Wiener space, we have an orthogonal basis of reproducing kernels associated with the sequence of integers, leading to what is known as the cardinal series or the Shannon sampling theorem.

Another way of defining de Branges spaces is as follows. We say that an entire function E belongs to the Hermite–Biehler class if it has no real zeros and satisfies $|E(z)| > |E(\bar{z})|$ for z in the upper half-plane. For a given function E in the Hermite–Biehler class, we let $H(E)$ be the Hilbert space consisting of all entire functions f such that both f/E and f^*/E belong to H^2 . Here $f^*(z) = \overline{f(\bar{z})}$ and H^2 is the Hardy space of the upper half-plane, viewed in the usual way as a subspace of $L^2(\mathbb{R})$. We set

$$\|f\|_{H(E)}^2 = \int_{-\infty}^{\infty} \frac{|f(x)|^2}{|E(x)|^2} dx.$$

Then $H(E)$ is a de Branges space, and every de Branges space can be obtained in this way via a function E in the Hermite–Biehler class. We arrive at the Paley–Wiener space by setting $E(z) = e^{-i\pi z}$.

It follows from the preceding remarks that Theorems 1.1 and 1.2 apply to de Branges spaces with orthogonal bases of reproducing kernels located at a sequence of nonzero real points γ_n such that $\inf_n |\gamma_{n+1}|/|\gamma_n| > 1$.

2.4. Second example: model subspaces of H^2

Given an inner function I in the upper half-plane, we define the *model subspace* K_I^2 as

$$K_I^2 = H^2 \ominus IH^2,$$

which is the orthogonal complement in H^2 of functions divisible by the inner function I . These spaces are, by a classical theorem of Beurling [8], the subspaces of H^2 that are invariant

with respect to the backward shift. We refer to [26–28] for information about the model theory related to the backward shift. The elements of K_I^2 (originally defined in \mathbb{C}^+) have meromorphic extensions into \mathbb{C} if the function I has such an extension. In this case, we have the relation $I = E^*/E$ and the map $f \mapsto f/E$ is a unitary map from $H(E)$ to K_I^2 . Thus de Branges spaces can be viewed as a subclass of the collection of all model subspaces of H^2 .

We now prove that every model subspace satisfies axiom (A2) from Subsection 2.1. This is obvious if we consider K_I^2 as a space of functions on the upper half-plane, but for our purposes it is essential that we also include those points on the real line at which point evaluation makes sense. We need the fact that the reproducing kernel for K_I^2 at some point ζ in the upper half-plane is

$$\kappa_\zeta(z) = \frac{i}{2\pi} \cdot \frac{1 - \overline{I(\zeta)}I(z)}{z - \bar{\zeta}}.$$

This formula extends to each point ζ on the real line at which every function in K_I^2 has a nontangential limit whose modulus is bounded by a constant times the H^2 -norm of the function. A paper of Ahern and Clark [1] gives that these are exactly the points ζ at which I has an angular derivative, that is, at which both I and I' have nontangential limits and $|I(\zeta)| = 1$.

LEMMA 2.2. *The Hilbert space K_I^2 , viewed as a space of functions on the set*

$$\Omega = \{z = x + iy : y \geq 0 \text{ and } f \mapsto f(z) \text{ is bounded}\},$$

satisfies axiom (A2) of Subsection 2.1.

To make the proof more transparent, we single out the main technical ingredient as a separate lemma.

LEMMA 2.3. *If x_0 is a point on the real line at which the point evaluation functional for K_I^2 is bounded, then $\|\kappa_{x_0+iy} - \kappa_{x_0}\|_{H^2} \rightarrow 0$ when $y \rightarrow 0$.*

Proof. Assuming $I(x_0) = 1$, we may write

$$\begin{aligned} \frac{2\pi}{i}(\kappa_{x_0+iy}(t) - \kappa_{x_0}(t)) &= \frac{1 - \overline{I(x_0 + iy)}I(t)}{t - x_0 + iy} - \frac{1 - I(t)}{t - x_0} \\ &= \frac{(1 - \overline{I(x_0 + iy)})I(t)}{t - x_0 + iy} - \frac{(1 - I(t))iyt}{(t - x_0)(t - x_0 + iy)}. \end{aligned}$$

Here the first term on the right-hand side has H^2 -norm bounded by a constant times $y^{1/2}$ in view of the theorem of Ahern and Clark [1], while the H^2 -norm of the second term tends to 0 when $y \rightarrow 0$, by Lebesgue's dominated convergence theorem. \square

Proof of Lemma 2.2. Let Λ be an exact uniqueness set for K_I^2 consisting of points in Ω . We let g_j denote the unique function in K_I^2 such that $g_j(\lambda_l)$ equals 0 when $l \neq j$ and 1 for $l = j$. We can choose an arbitrary point in Λ , say λ_1 , and choose $G(z) = (z - \lambda_1)g_1(z)$ as our candidate for a generating function. It is plain that if λ_j is a point in the open half-plane, then $g_j(z) = G(z)/[G'(\lambda_j)(z - \lambda_j)]$. The difficulty occurs if λ_j is a point on the real line. In this case, if we replace λ_j by $\lambda_j + i\varepsilon$, then the modified sequence $\Lambda^{(\varepsilon)}$ will still be an exact uniqueness sequence for K_I^2 . In fact, by Lemma 2.3, the function g_1 vanishing on $\Lambda^{(\varepsilon)} \setminus \{\lambda_1\}$ will vary continuously with ε . Thus the corresponding generating function $G_\varepsilon(z)$ will tend to $G(z)$ for every point in the upper half-plane when $\varepsilon \rightarrow 0$. On the other hand, another application of

Lemma 2.3 gives that $G_\varepsilon(z)/[G'_\varepsilon(\lambda_j + i\varepsilon)(z - \lambda_j + i\varepsilon)] \rightarrow g_j(z)$ in K_I^2 when $\varepsilon \rightarrow 0$. Lemma 2.3 also gives that $G'_\varepsilon(\lambda_j + i\varepsilon)$ converges to a finite number, say $1/\alpha$, and we may therefore conclude that $g_j(z) = \alpha G(z)/(z - \lambda_j)$. \square

As for axiom (A3), it remains an open problem to decide whether every model subspace K_I^2 has a Riesz basis of normalized reproducing kernels. Thus it is not known whether the class of spaces introduced in Subsection 2.1 includes all model subspaces. However, there exists an interesting class of model subspaces that actually possess orthogonal bases of reproducing kernels associated with sequences of real points. Such bases, to be discussed briefly below, are called Clark bases [16]. We also note that if the inner function I happens to be an interpolating Blaschke product, then it is well known and easy to show that K_I^2 has a Riesz basis of normalized reproducing kernels associated with the sequence of zeros of I .

The spaces K_I^2 that possess Clark bases correspond precisely to those spaces $\mathcal{H}(\Gamma, v)$ for which Γ is a real sequence. To get from $\mathcal{H}(\Gamma, v)$ to the corresponding space K_I^2 , we construct the Herglotz function

$$\varphi(z) = \sum_{n=1}^{\infty} v_n \left(\frac{1}{\gamma_n - z} - \frac{\gamma_n}{1 + \gamma_n^2} \right). \quad (2.5)$$

Then

$$I(z) = \frac{\varphi(z) - i}{\varphi(z) + i} \quad (2.6)$$

will be an inner function in the upper half-plane, and the map $f \mapsto (1 - I)f$ will be a unitary map from $\mathcal{H}(\Gamma, v)$ to K_I^2 ; it is implicit in this construction that in fact every function in K_I^2 has a nontangential limit at each point γ_n and also that the corresponding point evaluation functional is bounded at γ_n . We refer to our recent paper [7], where Clark bases are treated in more detail and it is shown that families of orthogonal bases of reproducing kernels can only exist in spaces of the form $\mathcal{H}(\Gamma, v)$ when Γ is a subset of a straight line or a circle.

We conclude that our general discussion (including Theorem 1.3) applies to model subspaces K_I^2 that possess Clark bases.

2.5. Carleson measures and systems of reproducing kernels in de Branges spaces and model subspaces of H^2

A long-standing problem in the function theory of de Branges spaces and, more generally, of model subspaces K_I^2 is to describe their Carleson measures. Only a few special cases have been completely understood. One such case is when I is a so-called one-component inner function, that is, when there exists a positive number ε with $0 < \varepsilon < 1$ such that the set of points z in the upper half-plane satisfying $|I(z)| < 1 - \varepsilon$ is connected [17, 30]. Other partial results can be found in [2–4, 18, 19, 30]. The lack of a general result on Carleson measures and, more specifically, on the geometry of Bessel sequences of normalized reproducing kernels is an obvious challenge when we address the Feichtinger conjecture in this setting.

Since positive results on Carleson measures are scarce, we mention without proof the following observation: a suitable adaption of Theorem 1.1 gives a description of any Carleson measure μ restricted to a cone $\{z = x + iy : |z - x_0| < Cy\}$; here x_0 is an arbitrary real point and C a positive constant. To arrive at this result, one may represent the space by means of its Clark basis or more generally as an L^2 -space with respect to a Clark measure [16], and act similarly as in Section 3.

A conjecture of Cohn [17], suggesting that it would suffice to verify the Carleson measure condition for reproducing kernels, was refuted by Nazarov and Volberg [25]. In the present paper, we likewise give a negative answer to the following question raised by Baranov.

QUESTION. Suppose that Γ and Λ are disjoint sequences of real numbers and that $\gamma_n \nearrow \infty$. If w is the Bessel weight sequence for Λ with respect to (Γ, v) and $H_{(\Gamma, v); (\Lambda, w)}$ is a bounded transformation, then is it true that there is a uniform bound on the number of λ_j found between two points γ_n and γ_{n+1} ?

A slight modification of our general approach will lead to a suitable example, to be presented in Subsection 3.4, with no such uniform bound.

It was recently shown by Baranov and Dyakonov [6] that the Feichtinger conjecture holds true for normalized reproducing kernels for K_I^2 when either I is a one-component inner function or the points λ_j associated with the sequence of reproducing kernels κ_{λ_j} satisfy

$$\sup_j |I(\lambda_j)| < 1. \quad (2.7)$$

In the latter case, a complete description of Riesz basic sequences exists [20], and this result plays an essential role in the proof. Baranov and Dyakonov used their result for the case when (2.7) holds to treat the general case of one-component inner functions. Their approach, following a technique used in [5], was to split the half-plane into two regions, one in which $|I(z)|$ is bounded away from 1 and another in which a perturbation argument for Clark bases applies.

Our results complement those of [6] and show that, in general, one needs additional ideas to resolve the Feichtinger conjecture for normalized reproducing kernels for model subspaces. First of all, since Γ satisfies the sparseness condition (1.2), the corresponding inner function I , defined by (2.6) via (2.5), is not a one-component inner function. Moreover, as will be revealed in Subsection 5.3, there exist inner functions I and Bessel sequences of normalized reproducing kernels $\kappa_{\lambda_j} / \|\kappa_{\lambda_j}\|_2$ for K_I^2 such that $|I(\lambda_j)| \rightarrow 1$ but no infinite subsequence is a subsequence of a Riesz basis.

By the observation made at the end of the previous subsection, the problem of describing all Riesz bases of normalized reproducing kernels for K_I^2 is part of the problem of deciding when discrete Hilbert transforms $H_{(\Gamma, v); (\Lambda, w)}$ are bounded. The most far-reaching result known about such bases is that found in [20] dealing with the case when (2.7) holds. The general result in [20] for this particular case leads to a description of all Riesz bases of normalized reproducing kernels for the Paley–Wiener space and also for a wider class of de Branges spaces known as weighted Paley–Wiener spaces [24]. One of the main points of [20] is that when (2.7) holds, one can transform the problem into a question about invertibility of Toeplitz operators and then apply the Devinatz–Widom theorem. Another approach, closer in spirit to the present work, can be found in [23], where the Riesz basis problem is explicitly related to a boundedness problem for Hilbert transforms.

2.6. Third example: ‘small’ Fock-type spaces

It may be noted that our work gives a full description of the Carleson measures and the Riesz bases of normalized reproducing kernels for certain ‘small’ Fock-type spaces studied recently by Borichev and Lyubarskii [9]. The spaces \mathcal{H} considered by these authors consist of all entire functions f such that

$$\|f\|_\varphi^2 = \int_{\mathbb{C}} |f(z)|^2 e^{-2\varphi(|z|)} dm(z) < \infty,$$

where φ is a positive, increasing, and unbounded function on $[0, \infty)$ and m denotes Lebesgue area measure on \mathbb{C} . The main point of [9] is that if φ grows ‘at most as fast’ as $[\log(1+r)]^2$, then the corresponding space \mathcal{H} has a Riesz basis of reproducing kernels and, conversely, if the growth of φ is ‘faster’ than $[\log(1+r)]^2$, then no such basis exists. It is proved that when $\varphi(r) = [\log(1+r)]^2$, we can choose such a basis associated with a sequence $\Gamma = (\gamma_n)$

satisfying $|\gamma_n| = e^{n/2}$; if $\varphi(r) = [\log(r+1)]^\alpha$ with $1 < \alpha < 2$, then the growth of $|\gamma_n|$ will be super-exponential.

In view of the observations in the previous subsections, the results of Borichev and Lyubarskii clarify when a Fock-type space equals a de Branges space, that is, when the two spaces consist of the same entire functions and have equivalent norms.

3. The boundedness problem

3.1. Proof of Theorem 1.1

In what follows, we use the notation

$$V_1 = 1, \quad V_n = \sum_{j=1}^{n-1} v_j \quad \text{and} \quad P_n = \sum_{j=n+1}^{\infty} \frac{v_j}{|\gamma_j|^2}. \quad (3.1)$$

Proof of the necessity of the conditions in Theorem 1.1. We observe first that the necessity of (1.4) is obvious: just apply $H_{(\Gamma, v)}$ to the sequence $e^{(n)} = (e_m^{(n)})$ with $e_n^{(n)} = 1$ and $e_m^{(n)} = 0$ for $m \neq n$.

To show that (1.5) is also a necessary condition, we begin by looking at the sequence $c^{(n)} = (c_m^{(n)})$ so that $c_m^{(n)} = 1$ for $m < n$ and $c_m^{(n)} = 0$ otherwise. We observe that $\|c^{(n)}\|_v^2 = V_n$ and note that, for z in Ω_l and $l \geq n$, we have

$$|H_{(\Gamma, v)} c^{(n)}(z)|^2 = \left| \sum_{m=1}^{n-1} \frac{v_m}{z - \gamma_m} \right|^2 \gtrsim \frac{V_n^2}{|z|^2}.$$

Taking into account the boundedness of $H_{(\Gamma, v)}$, we deduce from this that

$$V_n \gtrsim \int_{\mathbb{C}} |H_{(\Gamma, v)} c^{(n)}(z)|^2 d\mu(z) \gtrsim V_n^2 \sum_{m \geq n} \int_{\Omega_m} \frac{d\mu(z)}{|z|^2}.$$

On the other hand, if we set $a^{(n)} = (a_m^{(n)})$ so that $a_m^{(n)} = 1/\gamma_m$ for $m > n$ and $a_m^{(n)} = 0$ otherwise, then $\|a^{(n)}\|_v^2 = P_n$. We note that, for z in Ω_l and $l \leq n$, we have

$$|H_{(\Gamma, v)} a^{(n)}(z)|^2 = \left| \sum_{m=n+1}^{\infty} \frac{v_m}{\gamma_m(z - \gamma_m)} \right|^2 \gtrsim P_n^2.$$

Thus

$$P_n \gtrsim \int_{\mathbb{C}} |H_{(\Gamma, v)} a^{(n)}(z)|^2 d\mu(z) \gtrsim P_n^2 \sum_{m \leq n} \mu(\Omega_m). \quad \square$$

Proof of the sufficiency of the conditions in Theorem 1.1. Let $a = (a_n)$ be an arbitrary sequence in ℓ_v^2 . First we make the following estimate:

$$\begin{aligned} \int_{\Omega_n} |H_{(\Gamma, v)} a(z)|^2 d\mu(z) &\leq 3 \int_{\Omega_n} \left[\left| \sum_{m=1}^{n-1} \frac{a_m v_m}{z - \gamma_m} \right|^2 + \frac{|a_n|^2 v_n^2}{|z - \gamma_n|^2} + \left| \sum_{m=n+1}^{\infty} \frac{a_m v_m}{z - \gamma_m} \right|^2 \right] d\mu(z) \\ &\lesssim \int_{\Omega_n} \left[|z|^{-2} \left(\sum_{m=1}^{n-1} |a_m| v_m \right)^2 + \left(\sum_{m=n+1}^{\infty} \frac{|a_m| v_m}{|\gamma_m|} \right)^2 \right] d\mu(z) + |a_n|^2 v_n; \end{aligned}$$

here we used the Cauchy–Schwarz inequality and (1.4). Hence it remains for us to show that

$$\sum_{n=1}^{\infty} \left(\sum_{m=1}^{n-1} |a_m| v_m \right)^2 \int_{\Omega_n} |z|^{-2} d\mu(z) \lesssim \sum_{j=1}^{\infty} |a_j|^2 v_j \quad (3.2)$$

and

$$\sum_{n=1}^{\infty} \left(\sum_{m=n+1}^{\infty} \frac{|a_m| v_m}{|\gamma_m|} \right)^2 \mu(\Omega_n) \lesssim \sum_{j=1}^{\infty} |a_j|^2 v_j. \quad (3.3)$$

First we consider (3.2). To simplify the writing, we set

$$\tau_n = \left(\int_{\Omega_n} |z|^{-2} d\mu(z) \right)^{1/2}.$$

By duality,

$$\left(\sum_{n=1}^{\infty} \tau_n^2 \left(\sum_{m=1}^{n-1} |a_m| v_m \right)^2 \right)^{1/2} = \sup_{\|(c_n)\|_{\ell^2}=1} \sum_{n=1}^{\infty} |c_n| \tau_n \sum_{m=1}^{n-1} |a_m| v_m.$$

Since

$$\sum_{n=1}^{\infty} |c_n| \tau_n \sum_{m=1}^{n-1} |a_m| v_m = \sum_{m=1}^{\infty} |a_m| v_m \sum_{n=m+1}^{\infty} |c_n| \tau_n,$$

it suffices to show that the ℓ^2 -norm of

$$\alpha_m = v_m^{1/2} \sum_{n=m+1}^{\infty} |c_n| \tau_n$$

is bounded by a constant times the ℓ^2 -norm of (c_n) . To this end, we note that the Cauchy–Schwarz inequality gives

$$|\alpha_m|^2 \leq v_m \sum_{n=m+1}^{\infty} |c_n|^2 V_n^{-1/2} \sum_{j=m+1}^{\infty} \tau_j^2 V_j^{1/2}.$$

By (1.5), we see that

$$\sum_{j: 2^l V_m < V_j \leq 2^{l+1} V_m} \tau_j^2 V_j^{1/2} \lesssim \frac{1}{2^{l/2} V_{m+1}^{1/2}}$$

for $l \geq 0$. Summing these inequalities, we get

$$\sum_{j=m+1}^{\infty} \tau_j^2 V_j^{1/2} \lesssim \frac{1}{V_{m+1}^{1/2}}.$$

Hence

$$|\alpha_m|^2 \lesssim \frac{v_m}{V_{m+1}^{1/2}} \sum_{n=m+1}^{\infty} |c_n|^2 V_n^{-1/2}.$$

This gives us

$$\sum_{m=1}^{\infty} |\alpha_m|^2 \lesssim \sum_{n=1}^{\infty} |c_n|^2 V_n^{-1/2} \sum_{m=1}^{n-1} \frac{v_m}{V_{m+1}^{1/2}},$$

and so (3.2) follows because

$$V_n^{-1/2} \sum_{m=1}^{n-1} \frac{v_m}{V_{m+1}^{1/2}} \leq V_n^{-1/2} \int_0^{V_n} x^{-1/2} dx = 2.$$

We next consider (3.3). We note to begin with that the Cauchy–Schwarz inequality gives

$$\left(\sum_{m=n+1}^{\infty} \frac{|a_m|v_m}{|\gamma_m|} \right)^2 \leq \sum_{m=n+1}^{\infty} |a_m|^2 v_m P_{m-1}^{1/2} \sum_{j=n+1}^{\infty} \frac{v_j}{P_{j-1}^{1/2} |\gamma_j|^2}.$$

Since

$$\sum_{j=n+1}^{\infty} \frac{v_j}{P_{j-1}^{1/2} |\gamma_j|^2} \leq \int_0^{P_n} x^{-1/2} dx \leq 2P_n^{1/2},$$

it follows that

$$\sum_{n=1}^{\infty} \mu(\Omega_n) \left(\sum_{m=n+1}^{\infty} \frac{|a_m|v_m}{|\gamma_m|} \right)^2 \lesssim \sum_{n=1}^{\infty} \mu(\Omega_n) P_n^{1/2} \sum_{m=n+1}^{\infty} |a_m|^2 v_m P_m^{1/2},$$

which becomes

$$\sum_{n=1}^{\infty} \mu(\Omega_n) \left(\sum_{m=n+1}^{\infty} \frac{|a_m|v_m}{|\gamma_m|} \right)^2 \lesssim \sum_{m=1}^{\infty} |a_m|^2 v_m P_{m-1}^{1/2} \sum_{n=1}^{m-1} \mu(\Omega_n) P_n^{1/2}$$

when we change the order of summation. From (1.5) it follows that

$$\sum_{n=1}^{m-1} \mu(\Omega_n) P_n^{1/2} \lesssim \sum_{l=0}^{\infty} \sum_{n: 2^l P_{m-1} \leq P_n \leq 2^{l+1} P_{m-1}} \mu(\Omega_n) P_n^{1/2} \lesssim \frac{1}{P_{m-1}^{1/2}} \sum_{l=0}^{\infty} \frac{1}{2^{l/2}} \lesssim \frac{1}{P_{m-1}^{1/2}},$$

and we get (3.3). \square

3.2. Special cases

Condition (1.4) of Theorem 1.1 is a condition on the local behaviour of μ , while condition (1.5) deals with its global behaviour. Combining the two conditions, we see that (1.4) may be replaced by the stronger global condition

$$\sup_{n \geq 1} \int_{\mathbb{C}} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} < \infty.$$

We single out two cases in which (1.5) is automatically fulfilled once either this condition or the original one (1.4) holds.

COROLLARY 3.1. *Suppose that the sequence Γ satisfies the sparseness condition (1.2) and that the numbers v_n grow at least exponentially and that the numbers $v_n/|\gamma_n|^2$ decay at least exponentially with n . If μ is a nonnegative measure on \mathbb{C} with $\mu(\Gamma) = 0$, then the operator $H_{(\Gamma, v)}$ is bounded from ℓ_v^2 to $L^2(\mathbb{C}, \mu)$ if and only if*

$$\sup_{n \geq 1} \int_{\Omega_n} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} < \infty.$$

COROLLARY 3.2. *Suppose that the sequence Γ satisfies the sparseness condition (1.2) and that $\sum_n v_n < \infty$. If μ is a nonnegative measure on \mathbb{C} with $\mu(\Gamma) = 0$, then the operator $H_{(\Gamma, v)}$ is bounded from ℓ_v^2 to $L^2(\mathbb{C}, \mu)$ if and only if*

$$\sup_{n \geq 1} \int_{\mathbb{C}} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} < \infty.$$

Both corollaries follow immediately from Theorem 1.1.

3.3. Bessel sequences

We now switch to discrete Hilbert transforms of the form $H_{(\Gamma,v);(\Lambda,w)}$ with w the Bessel weight sequence for Λ with respect to (Γ, v) . As explained in Subsection 2.2, this means that we will be dealing with Bessel sequences for $\mathcal{H}(\Gamma, v)$.

We want to disentangle condition (1.5). To this end, we split any given sequence Λ into following three disjoint sequences:

$$\begin{aligned}\Lambda^{(0)} &= \left\{ \lambda \in \Lambda : \text{if } \lambda \text{ is in } \Omega_n, \text{ then } \frac{v_n}{|\lambda - \gamma_n|^2} \geq \max\left(\frac{V_n}{|\lambda|^2}, P_n\right) \right\}, \\ \Lambda^{(V)} &= \left\{ \lambda \in \Lambda : \text{if } \lambda \text{ is in } \Omega_n, \text{ then } \frac{V_n}{|\lambda|^2} > \max\left(\frac{v_n}{|\lambda - \gamma_n|^2}, P_n\right) \right\}, \\ \Lambda^{(P)} &= \left\{ \lambda \in \Lambda : \text{if } \lambda \text{ is in } \Omega_n, \text{ then } P_n > \max\left(\frac{v_n}{|\lambda - \gamma_n|^2}, \frac{V_n}{|\lambda|^2}\right) \right\}.\end{aligned}$$

We say that a sequence Λ is *V-lacunary* if

$$\sup_n \# \left(\Lambda \cap \bigcup_{m: 2^n \leq V_m \leq 2^{n+1}} \Omega_m \right) < \infty$$

and *P-lacunary* if

$$\sup_n \# \left(\Lambda \cap \bigcup_{m: 2^{-n-1} \leq P_m \leq 2^{-n}} \Omega_m \right) < \infty.$$

THEOREM 3.1. *Suppose the sequence Γ satisfies the sparseness condition (1.2) and that v is an admissible weight sequence for Γ . Let Λ be a sequence in $(\Gamma, v)^*$ and let w be the Bessel weight sequence for Λ with respect to (Γ, v) . Then $H_{(\Gamma,v);(\Lambda,w)}$ is a bounded transformation if and only if $\sup_n \#(\Lambda \cap \Omega_n) < \infty$, $\Lambda^{(V)}$ is a V-lacunary sequence, $\Lambda^{(P)}$ is a P-lacunary sequence, and*

$$\sup_{n \geq 1} \left(V_n \sum_{m \geq n} \sum_{\lambda \in \Lambda^{(0)} \cap \Omega_m} \frac{|\lambda - \gamma_m|^2}{v_m |\lambda|^2} + P_n \sum_{m \leq n} \sum_{\lambda \in \Lambda^{(0)} \cap \Omega_m} \frac{|\lambda - \gamma_m|^2}{v_m} \right) < \infty. \quad (3.4)$$

This splitting into a ‘super-thin’ sequence $\Lambda^{(V)} \cup \Lambda^{(P)}$ and a ‘distorted’ sequence $\Lambda^{(0)}$ represents a phenomenon not previously recorded, as far as we know. Corollaries 3.1 and 3.2, when restricted to the case of Bessel sequences, describe two situations in which the ‘super-thin’ part does not appear, for different reasons: Corollary 3.1 covers the case when V_n grows exponentially and P_n decays exponentially with n ; $\Lambda^{(V)}$ and $\Lambda^{(P)}$ can then both be ‘absorbed’ in $\Lambda^{(0)}$. Corollary 3.2 covers the case when V_n is uniformly bounded so that $\Lambda^{(V)}$ can only be a finite sequence; the sequence $\Lambda^{(P)}$ can again be ‘absorbed’ in $\Lambda^{(0)}$.

We conclude that the most interesting situation occurs when either $v_n/|\gamma_n|^2 = o(P_n)$ or $v_n = o(V_n)$ and $V_n \rightarrow \infty$ as $n \rightarrow \infty$. These two cases will be studied in depth in Section 5.

3.4. An example answering Baranov’s question

We now modify our construction to obtain an example that gives a negative answer to Baranov’s question posed in Subsection 2.5.

We assume that (t_n) is a sequence of positive numbers such that

$$\inf_{n \geq 1} \frac{t_{n+1}}{t_n} > 1.$$

In addition, we assume that, for each positive integer n , we have the following cluster of n points:

$$\gamma_{n,l} = t_n + l - 1, \quad 1 \leq l \leq n.$$

We denote this finite sequence by Γ_n and set

$$\Gamma = \bigcup_{n=1}^{\infty} \Gamma_n.$$

We consider the simplest case when the corresponding weight sequence v is identically 1, that is, $v_{n,l} = 1$ for every point $\gamma_{n,l}$ in Γ .

It may be noted that if we want to describe the measures μ for which $H_{(\Gamma,1)}$ is bounded from ℓ^2 to $L^2(\mathbb{C} \setminus \Gamma, \mu)$, then it suffices to consider the behaviour of μ in the Carleson squares

$$S_n = \{z = x + iy : |x - \gamma_{n,1}| \leq 2n, 0 \leq y \leq 4n\}.$$

Indeed, outside these squares, each cluster Λ_n has basically the same effect as if a single point were located at, say, $\lambda_{n,1}$ with weight n . This means that Theorem 1.1 applies to describe the behaviour of μ outside the squares S_n . In fact, by this observation, one may obtain a complete solution to the boundedness problem for these particular sequences Γ and v . We omit this description here and confine the discussion to a suitable example solving Baranov's problem.

The preceding remarks indicate that the sequence Λ should be placed inside the union of the squares S_n . We set

$$\lambda_{n,s} = \gamma_{n,1} - 2^s, \quad 0 \leq s \leq \log_2 n$$

and then $\Lambda_n = (\lambda_{n,s})_s$ with s running from 0 to $[\log_2 n]$ (the integer part of $\log_2 n$), and

$$\Lambda = \bigcup_{n=1}^{\infty} \Lambda_n.$$

We observe that we have

$$w_{n,s} = \left(\sum_{m=1}^{\infty} \sum_{l=1}^n \frac{1}{|\gamma_{m,l} - \lambda_{n,s}|^2} \right)^{-1} \simeq |\lambda_{n,s} - \gamma_{n,1}| = 2^s.$$

These numbers constitute the sequence w , which is the Bessel weight sequence for Λ with respect to (Γ, v) . We now make the following claim.

CLAIM. *If the sequences Γ , Λ , and w are constructed as above, then $H_{(\Gamma,1);(\Lambda,w)}$ is a bounded transformation.*

The interesting point, giving a negative answer to Baranov's question, is that there are more than $\log_2 n$ points from Λ between the neighbouring clusters Λ_{n-1} and Λ_n .

Proof of the Claim. Let $a = (a_{n,l})$ be an arbitrary ℓ^2 -sequence associated with Γ and set

$$H_{(\Gamma,1)}a(\lambda) = \sum_{n=1}^{\infty} \sum_{l=1}^n \frac{a_{n,l}}{\lambda - \gamma_{n,l}}.$$

An application of the Cauchy–Schwarz inequality gives

$$\sum_{s=0}^{[\log_2 n]} |H_{(\Gamma,1)}a(\lambda_{n,s})|^2 w_{n,s} \lesssim \frac{n^3}{t_n^2} \|a\|_{\ell^2}^2 + \sum_{s=0}^{[\log_2 n]} 2^s \left(\sum_{l=1}^n \frac{|a_{n,l}|}{|\lambda_{n,s} - \gamma_{n,l}|} \right)^2.$$

The summation over n of the first term on the right-hand side causes no problem because t_n grows at least exponentially with respect to n . We therefore concentrate on the second term

$$A_n = \sum_{s=0}^{\lfloor \log_2 n \rfloor} 2^s \left(\sum_{l=1}^n \frac{|a_{n,l}|}{2^s + l - 1} \right)^2.$$

The Cauchy–Schwarz inequality gives

$$\left(\sum_{l=1}^n \frac{|a_{n,l}|}{2^s + l - 1} \right)^2 \leq \sum_{j=1}^n \frac{j^{-1/2}}{2^s + j - 1} \sum_{l=1}^n \frac{l^{1/2} |a_{n,l}|^2}{2^s + l - 1}.$$

The first of the two sums on the right-hand side is bounded by a constant times $2^{-s/2}$, and so it follows that

$$A_n \leq \sum_{s=0}^{\lfloor \log_2 n \rfloor} 2^{s/2} \sum_{l=1}^n \frac{l^{1/2} |a_{n,l}|^2}{2^s + l - 1}.$$

Changing the order of summation and using that

$$\sum_{s=0}^{\lfloor \log_2 n \rfloor} \frac{2^{s/2}}{2^s + l - 1} \lesssim l^{-1/2},$$

we finally obtain the desired estimate

$$A_n \lesssim \sum_{j=1}^n |a_{n,j}|^2. \quad \square$$

4. Proof of Theorem 1.2

We use the same splitting as in Theorem 3.1 and treat the three sequences $\Lambda^{(0)}$, $\Lambda^{(V)}$, and $\Lambda^{(P)}$ separately. We use Lemma 2.1, that is, we make a splitting so that, for each subsequence Λ' with associated weight sequence, the adjoint transformation $H_{(\Lambda', w'); (\Gamma, v)}$ is bounded below. From now on, we shall use the notation

$$W_n = \sum_{m=1}^{n-1} w_m \quad \text{and} \quad Q_n = \sum_{m=n+1}^{\infty} \frac{w_m}{|\lambda_m|^2}. \quad (4.1)$$

4.1. The splitting of $\Lambda^{(0)}$

We may assume that there is at most one point λ_n in $\Lambda^{(0)}$ from each annulus Ω_n ; we denote the corresponding weights by w_n . Let $\Lambda' = (\lambda_{n_j})$ be a subsequence of $\Lambda^{(0)}$ with corresponding weight sequence $w' = (w_{n_j})$, and let $a = (a_{n_j})$ be an arbitrary $\ell_{w', v}^2$ -sequence. Since

$$|\xi - \eta|^2 \geq |\xi|^2 - 2|\xi||\eta| + |\eta|^2 \geq \frac{1}{2}|\xi|^2 - |\eta|^2$$

for arbitrary complex numbers ξ and η , we have

$$|H_{(\Lambda', w'); (\Gamma, v)} a(\gamma_{n_j})|^2 \geq \frac{1}{2} \frac{|a_{n_j}|^2 w_{n_j}^2}{|\lambda_{n_j} - \gamma_{n_j}|^2} - 2 \left| \sum_{l=1}^{j-1} \frac{a_{n_l} w_{n_l}}{\gamma_{n_j} - \lambda_{n_l}} \right|^2 - 2 \left| \sum_{l=j+1}^{\infty} \frac{a_{n_l} w_{n_l}}{\gamma_{n_j} - \lambda_{n_l}} \right|^2.$$

On the other hand, $w_{n_j} \simeq |\lambda_{n_j} - \gamma_{n_j}|^2 / v_{n_j}$. Therefore, by the definition of $\Lambda^{(0)}$, there is a positive constant c such that

$$\|H_{(\Lambda', w'); (\Gamma, v)} a\|_v^2 \geq c \|a\|_{w'}^2 - 2 \sum_{j=1}^{\infty} \left[\left| \sum_{l=1}^{j-1} \frac{a_{n_l} w_{n_l}}{\bar{\gamma}_{n_j} - \bar{\lambda}_{n_l}} \right|^2 + \left| \sum_{l=j+1}^{\infty} \frac{a_{n_l} w_{n_l}}{\bar{\gamma}_{n_j} - \bar{\lambda}_{n_l}} \right|^2 \right] v_{n_j}. \quad (4.2)$$

Hence it remains for us to show that, for a given $\varepsilon > 0$, we may obtain

$$\sum_{j=1}^{\infty} \left(\sum_{l=1}^{j-1} |a_{n_l} w_{n_l}| \right)^2 \frac{v_{n_j}}{|\lambda_{n_j}|^2} \leq \varepsilon \sum_{j=1}^{\infty} |a_{n_j}|^2 w_{n_j} \quad (4.3)$$

and

$$\sum_{j=1}^{\infty} \left(\sum_{l=j+1}^{\infty} \frac{|a_{n_l} w_{n_l}|}{|\lambda_{n_l}|} \right)^2 v_{n_j} \leq \varepsilon \sum_{j=1}^{\infty} |a_{n_j}|^2 w_{n_j} \quad (4.4)$$

for every subsequence Λ' in a finite splitting of $\Lambda^{(0)}$.

We proceed as in the proof of Theorem 1.1. Thus we set $\tau_j = v_{n_j}^{1/2} / |\lambda_{n_j}|$ and consider first (4.3). By duality,

$$\left(\sum_{j=1}^{\infty} \tau_j^2 \left(\sum_{l=1}^{j-1} |a_{n_l} w_{n_l}| \right)^2 \right)^{1/2} = \sup_{\|(c_j)\|_{\ell^2}=1} \sum_{j=1}^{\infty} |c_j| \tau_j \sum_{l=1}^{j-1} |a_{n_l} w_{n_l}|.$$

Since

$$\sum_{j=1}^{\infty} |c_j| \tau_j \sum_{l=1}^{j-1} |a_{n_l} w_{n_l}| = \sum_{l=1}^{\infty} |a_{n_l} w_{n_l}| \sum_{j=l+1}^{\infty} |c_j| \tau_j,$$

it suffices to show that the ℓ^2 -norm of

$$\alpha_l = w_{n_l}^{1/2} \sum_{j=l+1}^{\infty} |c_j| \tau_j$$

can be made smaller than ε times the ℓ^2 -norm of (c_j) . To this end, we note that the Cauchy-Schwarz inequality gives

$$|\alpha_l|^2 \leq w_{n_l} \sum_{j=l+1}^{\infty} |c_j|^2 W_{n_j}^{-1/2} \sum_{m=l+1}^{\infty} \tau_m^2 W_{n_m}^{1/2}.$$

Using (1.5), we get

$$\sum_{m=l+1}^{\infty} \tau_m^2 W_{n_m}^{1/2} \lesssim \frac{1}{W_{n_{l+1}}^{1/2}}.$$

Hence

$$|\alpha_l|^2 \lesssim \frac{w_{n_l}}{W_{n_{l+1}}^{1/2}} \sum_{j=l+1}^{\infty} |c_j|^2 W_{n_j}^{-1/2}.$$

This gives us

$$\sum_{l=1}^{\infty} |\alpha_l|^2 \lesssim \sum_{j=1}^{\infty} |c_j|^2 W_{n_j}^{-1/2} \sum_{l=1}^{j-1} \frac{w_{n_l}}{W_{n_{l+1}}^{1/2}},$$

and so (4.3) would follow if we could obtain

$$\sum_{l=1}^{j-1} \frac{w_{n_l}}{W_{n_{l+1}}^{1/2}} \leq c\varepsilon W_{n_j}^{1/2} \quad (4.5)$$

for an absolute constant c .

Having singled out this goal, we proceed to consider (4.4). We note to begin with that the Cauchy–Schwarz inequality gives

$$\left(\sum_{l=j+1}^{\infty} \frac{|a_{n_l}| w_{n_l}}{|\lambda_{n_j}|} \right)^2 \leq \sum_{l=j+1}^{\infty} |a_{n_l}|^2 w_{n_l} Q_{n_{l-1}}^{1/2} \sum_{m=j+1}^{\infty} \frac{w_{n_m}}{Q_{n_{m-1}}^{1/2} |\lambda_{n_m}|^2}.$$

Now our goal will be to obtain

$$\sum_{m=j+1}^{\infty} \frac{w_{n_m}}{Q_{n_{m-1}}^{1/2} |\lambda_{n_m}|^2} \leq c\varepsilon Q_{n_j}^{1/2}. \quad (4.6)$$

Indeed, this would imply

$$\begin{aligned} \sum_{j=1}^{\infty} \left(\sum_{l=j+1}^{\infty} \frac{|a_{n_l}| w_{n_l}}{|\lambda_{n_j}|} \right)^2 v_{n_j} &\lesssim \varepsilon \sum_{j=1}^{\infty} v_{n_j} Q_{n_j}^{1/2} \sum_{l=j+1}^{\infty} |a_{n_l}|^2 w_{n_l} Q_{n_{l-1}}^{1/2} \\ &= \varepsilon \sum_{l=1}^{\infty} |a_{n_l}|^2 w_{n_l} Q_{n_{l-1}}^{1/2} \sum_{j=1}^{l-1} v_{n_j} Q_{n_j}^{1/2}. \end{aligned}$$

By (1.5), we have

$$\sum_{j=1}^{l-1} v_{n_j} Q_{n_j}^{1/2} \lesssim \frac{1}{Q_{n_{l-1}}^{1/2}},$$

and so it will suffice to have (4.6).

In order to obtain the two estimates (4.5) and (4.6) for every subsequence in our finite splitting of $\Lambda^{(0)}$, we make a splitting according to the following algorithm:

(1) Let δ be a small positive number to be chosen later. Select those n for which $w_n > \delta W_n$. If we choose Λ' to consist of every N th λ_n in the corresponding subsequence of $\Lambda^{(0)}$, then we get

$$\sum_{l=1}^{j-1} \frac{w_{n_l}}{W_{n_{l+1}}^{1/2}} \leq \frac{2}{\delta N} W_{n_j}^{1/2}$$

by again comparing the sum to the integral of the function $x^{-1/2}$ over the interval from 0 to W_{n_j} . Thus we achieve our goal if we choose N to be of the order of magnitude $1/(\delta\varepsilon)$.

(2) Return to those points λ_{n_j} not selected in (1). For these we have $w_{n_j} \leq \delta W_{n_j}$. Group these points into blocks of points with consecutive indices such that, for each block, $\delta \leq \sum w_{n_j} W_{n_j}^{-1} < 2\delta$. Construct new subsequences by picking every N th block from this sequence of blocks. Then some elementary estimates, again using comparisons with an integral, lead to the following inequality:

$$\sum_{l=1}^{j-1} \frac{w_{n_l}}{W_{n_{l+1}}^{1/2}} \leq \frac{16\delta}{1 - (1 - 2\delta)^N} W_{n_j}^{1/2},$$

where we sum over the new subsequence. Thus it would suffice if we choose N to be roughly $1/\delta$ and δ to be a suitable constant times ε .

(3) Take one of the subsequences selected in (1) or (2) and consider the subsequence of this subsequence, say $\Lambda' = (\lambda_{n_j})$, along which $w_{n_j} |\lambda_{n_j}|^{-2} > \delta Q_{n_j}$. If we select a new subsequence

by picking every N th λ_{n_j} in the sequence Λ' , then the sum in (4.6) becomes smaller than $2/(\delta N)Q_{n_j}$ by the same argument as in (1). Again our goal is achieved if we choose N to be of the order of magnitude $1/(\delta\varepsilon)$.

(4) Take again one of the subsequences selected in (1) or (2) and consider those subsequences of these for which we have $w_{n_j}|\lambda_{n_j}|^{-2} \leq \delta Q_{n_j}$. Group the points in these subsequences into blocks of points with consecutive indices such that for each block $\delta \leq \sum w_{n_j}|\lambda_{n_j}|^{-2} Q_{n_j}^{-1} < 2\delta$. Now construct new subsequences by picking every N th block from this sequence of blocks. Then as in point (2) we get

$$\sum_{m=j+1}^{\infty} \frac{w_{n_m}}{Q_{n_{m-1}}^{1/2} |\lambda_{n_m}|^2} \leq \frac{16\delta}{1 - (1 - 2\delta)^N} Q_{n_j}^{1/2}.$$

(Here the summation is again over the new subsequence.) We observe once more that it would suffice if we choose N to be roughly $1/\delta$ and δ to be a suitable constant times ε .

4.2. The splitting of $\Lambda^{(V)}$

The splitting of $\Lambda^{(V)}$ is almost identical to that of $\Lambda^{(0)}$. We now use the estimate

$$|H_{(\Lambda', w'); (\Gamma, v)} a(\gamma_n)|^2 \geq \frac{1}{2} \frac{|a_{n_j}|^2 w_{n_j}^2}{|\lambda_{n_j} - \gamma_n|^2} - 2 \left| \sum_{l=1}^{j-1} \frac{a_{n_l} w_{n_l}}{\lambda_{n_j} - \gamma_n} \right|^2 - 2 \left| \sum_{l=j+1}^{\infty} \frac{a_{n_l} w_{n_l}}{\gamma_n - \lambda_{n_l}} \right|^2. \quad (4.7)$$

The reason we write ' γ_n ' instead of ' γ_{n_j} ' is that we need to sum over several annuli Ω_n in order to estimate the norm of $\|a\|_w$. Indeed, we may assume that λ_{n_j} belongs to a union of annuli Ω_n , denoted by Δ_j , such that

$$\sum_{\gamma_n \in \Delta_j} \frac{v_n}{|\lambda_{n_j} - \gamma_n|^2} \geq \frac{1}{10} \frac{V_{n_j}}{|\lambda_j|^2},$$

with the sets Δ_j being pairwise disjoint. Therefore, by the definition of $\Lambda^{(V)}$, there is a constant c such that

$$\sum_{\gamma_n \in \Delta_j} |a_{n_j}|^2 w_{n_j}^2 \frac{v_n}{|\lambda_{n_j} - \gamma_n|^2} \geq c |a_{n_j}|^2 w_{n_j}.$$

Hence we obtain

$$\|H_{(\Lambda', w'); (\Gamma, v)} a\|_v^2 \geq c \|a\|_{w'}^2 - 2 \sum_{j=1}^{\infty} \sum_{\gamma_n \in \Delta_j} \left[\left| \sum_{l=1}^{j-1} \frac{a_{n_l} w_{n_l}}{\lambda_{n_j} - \gamma_n} \right|^2 + \left| \sum_{l=j+1}^{\infty} \frac{a_{n_l} w_{n_l}}{\gamma_n - \lambda_{n_l}} \right|^2 \right] v_{n_j}.$$

The splitting is then done in essentially the same way as above, repeating the reasoning based on the estimate (4.2).

4.3. The splitting of $\Lambda^{(P)}$

We use once more (4.7), but this time we may assume that λ_{n_j} belongs to a union of annuli Ω_n , again denoted by Δ_j , such that

$$\sum_{\gamma_n \in \Delta_j} \frac{v_n}{|\lambda_{n_j} - \gamma_n|^2} \geq \frac{1}{10} P_{n_j},$$

with the sets Δ_j being pairwise disjoint. Therefore, by the definition of $\Lambda^{(P)}$, there is a constant c such that

$$\sum_{\gamma_n \in \Delta_j} |a_{n_j}|^2 w_{n_j}^2 \frac{v_n}{|\lambda_{n_j} - \gamma_n|^2} \geq c |a_{n_j}|^2 w_{n_j}.$$

Hence we obtain

$$\|H_{(\Lambda', w'); (\Gamma, v)} a\|_v^2 \geq c \|a\|_{w'}^2 - 2 \sum_{j=1}^{\infty} \sum_{\gamma_n \in \Delta_j} \left[\left| \sum_{l=1}^{j-1} \frac{a_{n_l} w_{n_l}}{\lambda_{n_j} - \gamma_n} \right|^2 + \left| \sum_{l=j+1}^{\infty} \frac{a_{n_l} w_{n_l}}{\lambda_{n_j} - \gamma_n} \right|^2 \right] v_{n_j},$$

and proceed as outlined in the previous paragraph.

5. The invertibility problem

5.1. Proof of Theorem 1.3

It is clear that if the mapping $H_{(\Gamma, v); (\Lambda, w)}$ is invertible; then Λ is an exact uniqueness sequence for $H_{(\Gamma, v)}$, which in turn implies that there is a unique element $e = (e_n)$ in ℓ_v^2 such that $H_{(\Gamma, v)} e$ vanishes on $\Gamma \setminus \{\lambda_1\}$ and takes the value 1 at λ_1 . Then

$$G(z) = (z - \lambda_1) \sum_{n=1}^{\infty} \frac{e_n v_n}{z - \gamma_n}$$

is a generating function for Λ . Since by assumption $G(\lambda_j) = 0$ for $j > 1$, we may write

$$G(z) = G(z) - G(\lambda_j) = (z - \lambda_j) \sum_{n=1}^{\infty} \frac{e_n v_n (\gamma_n - \lambda_1)}{(\gamma_n - \lambda_j)(z - \gamma_n)},$$

where on the right-hand side we have just subtracted the respective series that define $G(z)$ and $G(\lambda_j)$. Since G is a generating function for Λ , it follows that

$$\sum_{n=1}^{\infty} \frac{|e_n|^2 |\gamma_n - \lambda_1|^2 v_n}{|\gamma_n - \lambda_j|^2} < \infty.$$

In particular, the sequence

$$e^{(j)} = \left(e_n \frac{\gamma_n - \lambda_1}{\gamma_n - \lambda_j} \left(\sum_{m=1}^{\infty} \frac{e_m v_m (\lambda_1 - \gamma_m)}{(\lambda_j - \gamma_m)^2} \right)^{-1} \right)_n$$

will be the unique vector in ℓ_v^2 such that $H_{(\Gamma, v)} e^{(j)}(\lambda_l)$ is 0 when $l \neq j$ and 1 for $l = j$.

To simplify the writing, we set

$$\alpha_j = \left(\sum_{m=1}^{\infty} \frac{e_m v_m (\lambda_1 - \gamma_m)}{(\lambda_j - \gamma_m)^2} \right)^{-1};$$

thus if $b = (b_1, b_2, \dots, b_l, 0, 0, \dots)$ is a sequence with only finitely many nonzero entries, then the sequence

$$a = \left(e_n (\gamma_n - \lambda_1) \sum_{j=1}^l \frac{b_j \alpha_j}{\gamma_n - \lambda_j} \right)_n \quad (5.1)$$

will be the unique vector in ℓ_v^2 such that $H_{(\Gamma, v); (\Lambda, w)} a = b$. This means that we have identified a linear transformation, defined on a dense subset of ℓ_w^2 , that must be the inverse transformation to $H_{(\Gamma, v); (\Lambda, w)}$, should it exist. Hence, under the assumption that Λ is an exact uniqueness sequence for $H_{(\Gamma, v)}$, a necessary and sufficient condition for invertibility of $H_{(\Gamma, v); (\Lambda, w)}$ is that the linear transformation defined by (5.1) extends to a bounded transformation on ℓ_w^2 . An equivalent condition is that the transformation $H_{(\Lambda, \varpi); (\Gamma, \nu)}$ be bounded where

$$\nu_n = v_n |\lambda_1 - \gamma_n|^2 |e_n|^2$$

and

$$\varpi_j = w_j^{-1} \left| \sum_{n=1}^{\infty} \frac{e_n v_n (\lambda_1 - \gamma_n)}{(\lambda_j - \gamma_n)^2} \right|^{-2} = w_j^{-1} |\lambda_j - \lambda_1|^{-2} \left| \sum_{n=1}^{\infty} \frac{e_n v_n}{(\lambda_j - \gamma_n)^2} \right|^{-2}.$$

In the final step, we used the definition of the sequence (e_n) .

5.2. Localization of Λ when Γ is a sparse sequence

We will for the rest of this section consider two interesting special cases. The main point of this subsection will be that, although Λ may possibly have a nontrivial splitting into three sequences $\Lambda^{(0)}$, $\Lambda^{(V)}$, and $\Lambda^{(P)}$ (cf. the discussion in Subsection 3.3), the invertibility of $H_{(\Gamma, v); (\Lambda, w)}$ forces the sequences $\Lambda^{(V)}$ and $\Lambda^{(P)}$ to be trivial, in a sense to be made precise.

We assume as before that $\Gamma = (\gamma_n)$ is indexed by the positive integers, and that the sequence is sparse in the sense that (1.2) holds. We retain the notation

$$V_n = \sum_{m=1}^{n-1} v_m \quad \text{and} \quad P_n = \sum_{m=n+1}^{\infty} \frac{v_m}{|\gamma_m|^2}$$

from the previous section. In the discussion below, the sets

$$D_n(v; M) = \left\{ \lambda \in \Omega_n : \frac{M v_n}{|\lambda - \gamma_n|^2} \geq \max \left(\frac{V_n}{|\lambda|^2}, P_n \right) \right\},$$

defined for every admissible weight sequence v and positive number M , will play an essential role. If M is fixed and either $v_n = o(V_n)$ or $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$, then these sets are essentially discs centred at γ_n with radii that are $o(|\gamma_n|)$ when $n \rightarrow \infty$. In such situations, the splitting of a sequence Λ into the three sequences $\Lambda^{(0)}$, $\Lambda^{(V)}$, and $\Lambda^{(P)}$ may be nontrivial, in the sense that $\Lambda \setminus \bigcup_n D_n(v; M)$ may be an infinite sequence for every positive M .

We assume that $\Lambda = (\lambda_n)$ is a sequence disjoint from Γ , indexed by a sequence of integers $(n_0, n_0 + 1, n_0 + 2, \dots)$ and ordered such that the moduli $|\lambda_n|$ increase with n . For convenience, we assume that $\lambda_{n_0} \neq 0$. The choice of n_0 is made such that Λ is ‘aligned’ with Γ . More precisely, we say that Λ is a v -perturbation of Γ if n_0 can be chosen such that, for a sufficiently large M , λ_n is in $D_n(v; M)$ for all but possibly a finite number of indices n . If Λ is a v -perturbation of Γ , it will be implicitly understood that n_0 is chosen so that the two sequences are ‘aligned’ in this way.

A v -perturbation Λ of Γ will be said to be

- (i) an *exact* v -perturbation of Γ if $n_0 = 1$;
- (ii) a v -perturbation of Γ of *deficiency* $n_0 - 1$ if $n_0 > 1$;
- (iii) a v -perturbation of Γ of *excess* $1 - n_0$ if $n_0 < 1$.

The main results of this subsection are the following two lemmas.

LEMMA 5.1. *Suppose that w is the Bessel weight sequence for Λ with respect to (Γ, v) and that $v_n = o(V_n)$ when $n \rightarrow \infty$. If, in addition, the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is invertible, then Λ is either an exact v -perturbation of Γ or a v -perturbation of deficiency 1.*

LEMMA 5.2. *Suppose that w is the Bessel weight sequence for Λ with respect to (Γ, v) and that $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$. If, in addition, the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is invertible, then Λ is either an exact v -perturbation of Γ or a v -perturbation of Γ of excess 1.*

Note the contrast between these results and Theorem 3.1; Λ has no nontrivial V -lacunary or P -lacunary subsequences when $H_{(\Gamma, v); (\Lambda, w)}$ is an invertible transformation. We see in the next section that, quite remarkably, all three cases, exactness, deficiency 1, and excess 1, may occur.

The proofs of the two lemmas require several steps. We begin with a simple estimate, to be used repeatedly in what follows. It concerns the quantity

$$\varrho_n = \prod_{m=\max(1, n_0)}^n \frac{|\gamma_m|^2}{|\lambda_m|^2},$$

which will appear prominently in our conditions for invertibility. We use again the notation introduced in (4.1), that is, we set

$$W_n = \sum_{m=n_0}^{n-1} w_m \quad \text{and} \quad Q_n = \sum_{m=n+1}^{\infty} \frac{w_m}{|\lambda_m|^2}.$$

LEMMA 5.3. *If Λ is a v -perturbation of Γ and $|\gamma_n| \simeq |\lambda_n|$, then we have both*

$$\left| \log \frac{\varrho_m}{\varrho_n} \right|^2 \lesssim (V_{m+1} - V_{n+1})(Q_n - Q_m) \quad (5.2)$$

and

$$\left| \log \frac{\varrho_m}{\varrho_n} \right|^2 \lesssim (W_{m+1} - W_{n+1})(P_n - P_m) \quad (5.3)$$

when $m > n$. If, in addition, either $v_n = o(V_n)$ or $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$, then $\log \varrho_n = o(n)$ when $n \rightarrow \infty$.

Proof. Since $|\gamma_n| \simeq |\lambda_n|$, we have

$$\left| \log \frac{\varrho_m}{\varrho_n} \right| = 2 \left| \sum_{l=n+1}^m \log \frac{|\gamma_l|}{|\lambda_l|} \right| \lesssim \sum_{l=n+1}^m \left| 1 - \frac{|\gamma_l|}{|\lambda_l|} \right|. \quad (5.4)$$

Hence, by the Cauchy–Schwarz inequality, we get

$$\left| \log \frac{\varrho_m}{\varrho_n} \right|^2 \lesssim \sum_{l=n+1}^m v_l \sum_{j=n+1}^m \frac{|\gamma_j - \lambda_j|^2}{v_j |\lambda_j|^2},$$

which is the desired estimate (5.2). Another application of the Cauchy–Schwarz inequality to (5.4) gives

$$\left| \log \frac{\varrho_m}{\varrho_n} \right|^2 \lesssim \sum_{l=n+1}^m \frac{|\gamma_l - \lambda_l|^2}{v_l} \sum_{j=n+1}^m \frac{v_j}{|\gamma_j|^2},$$

which is the second estimate (5.3).

Finally, starting again from (5.4) and using the Cauchy–Schwarz inequality a third time, we get

$$|\log \varrho_n|^2 \lesssim n \sum_{l=\max(1, n_0)}^n \frac{|\gamma_l - \lambda_l|^2}{|\lambda_l|^2} \lesssim n \sum_{l=\max(1, n_0)}^n \min \left(\frac{v_l}{V_l}, \frac{v_l}{|\gamma_l|^2 P_l} \right),$$

where in the last step we used that Λ is a v -perturbation of Γ . This relation gives the last statement in the lemma, namely that $\log \varrho_n = o(n)$ when either $v_n = o(V_n)$ or $v_n/|\gamma_n|^2 = o(P_n)$ as $n \rightarrow \infty$. \square

We next prove the following lemma, which is really a corollary to Theorem 1.1.

LEMMA 5.4. *Suppose that either $v_n = o(V_n)$ or $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$. If, in addition, μ is a nonnegative measure on \mathbb{C} with $\mu(\Gamma) = 0$ and the map $H_{(\Gamma,v)}$ is both bounded and bounded below from ℓ_v^2 to $L^2(\mathbb{C}, \mu)$, then there exist positive numbers M and δ such that*

$$\int_{D_n(v;M)} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} \geq \delta$$

for all but finitely many indices n .

Proof. Applying the assumption about boundedness below to any sequence with only one nonzero entry, we find that there is a positive number σ independent of n such that

$$\int_{\mathbb{C}} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} \geq \sigma$$

for every n . On the other hand, since $|\gamma_n|$ grows at least exponentially and $H_{(\Gamma,v)}$ is bounded from ℓ_v^2 to $L^2(\mathbb{C}, \mu)$, we have

$$\sum_{m=n+1}^{\infty} \int_{\Omega_m} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} \lesssim v_n \sum_{m=n+1}^{\infty} \int_{\Omega_m} \frac{d\mu(z)}{|z|^2} \lesssim \min \left(\frac{v_n}{V_n}, \frac{v_n}{|\gamma_n|^2 P_n} \right)$$

and

$$\sum_{m=1}^{n-1} \int_{\Omega_m} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} \lesssim \frac{v_n}{|\gamma_n|^2} \sum_{m=1}^{n-1} \int_{\Omega_m} d\mu(z) \lesssim \min \left(\frac{v_n}{V_n}, \frac{v_n}{|\gamma_n|^2 P_n} \right).$$

We also have

$$\int_{\Omega_n \setminus D_n(v;M)} \frac{v_n d\mu(z)}{|z - \gamma_n|^2} \leq \frac{1}{M} \int_{\Omega_n} \max \left(\frac{V_n}{|\lambda|^2}, P_n \right) d\mu(z) \lesssim \frac{1}{M},$$

again using the condition for boundedness of the map $H_{(\Gamma,v)} : \ell_v^2 \rightarrow L^2(\mathbb{C}, \mu)$. The result follows with $\delta = \sigma/2$ if we choose a sufficiently large M . \square

The preceding lemma shows that if the transformation $H_{(\Gamma,v);(\Lambda,w)}$ is invertible, then Λ must contain a subsequence that is, a v -perturbation of Γ . The next two lemmas show that Λ itself must be a v -perturbation of Γ .

LEMMA 5.5. *Suppose that $v_n = o(V_n)$ when $n \rightarrow \infty$. If, in addition, Λ is an exact v -perturbation of Γ , then Λ is a uniqueness sequence for $H_{(\Gamma,v)}$.*

Proof. We argue by contradiction. So suppose that there is a nonzero vector $a = (a_n)$ in ℓ_v^2 such that $H_{(\Gamma,v)}a$ vanishes on Λ . This means that there is a nonzero entire function $J(z)$ such that

$$\sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} = J(z) \prod_{m=1}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_m}$$

for every z in $\mathbb{C} \setminus \Gamma$. If we now choose M sufficiently large, then we have

$$\frac{V_n}{|z|^2} \gtrsim |J(z)|^2 \varrho_n$$

for z in $\Omega_n \setminus D_n(v;M)$. Since $v_n = o(V_n)$ when $n \rightarrow \infty$, the left-hand side is bounded by $e^{-\delta n}$ for some positive δ , while, by Lemma 5.3, $\varrho_n = e^{o(n)}$ when $n \rightarrow \infty$. Thus the maximum of $|J(z)|$ in $\Omega_n \setminus D_n(v;M)$ tends to 0 when $n \rightarrow \infty$, which is a contradiction unless $J(z) \equiv 0$. \square

LEMMA 5.6. *Suppose that $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$. If, in addition, Λ is a v -perturbation of Γ of excess 1, then Λ is a uniqueness sequence for $H_{(\Gamma,v)}$.*

Proof. We argue again by contradiction and assume that there is a nonzero vector $a = (a_n)$ in ℓ_v^2 such that $H_{(\Gamma,v)}a$ vanishes on Λ . In this case, it follows that there is a nonzero entire function $J(z)$ such that

$$\sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} = J(z)(z - \lambda_0) \prod_{m=1}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_m}$$

for every z in $\mathbb{C} \setminus \Gamma$. If we now choose M sufficiently large, then we have

$$P_n \gtrsim |J(z)|^2 |z|^2 \varrho_n$$

for z in $\Omega_n \setminus D_n(v; M)$. Since $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$, we have that $P_n/|z|^2$ is bounded by $e^{-\delta n}$ for some positive number δ , while, by Lemma 5.3, $\varrho_n = e^{o(n)}$ when $n \rightarrow \infty$. Thus the maximum of $|J(z)|$ in $\Omega_n \setminus D_n(v; M)$ tends to 0 when $n \rightarrow \infty$, which is a contradiction unless $J(z) \equiv 0$. \square

We finally prove two lemmas that, together with the previous three lemmas, give the precise restrictions stated in Lemmas 5.1 and 5.2.

LEMMA 5.7. *Suppose that $v_n = o(V_n)$ when $n \rightarrow \infty$. If, in addition, Λ is a v -perturbation of Γ of deficiency 2, then Λ is not a uniqueness sequence for $H_{(\Gamma,v)}$.*

Proof. We may write

$$\frac{c}{(z - \gamma_1)(z - \gamma_2)} \prod_{m=3}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_n} = \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} + h(z),$$

where h is an entire function and

$$|a_n|^2 v_n^2 \simeq \frac{|\gamma_n - \lambda_n|^2}{|\gamma_n|^4} \varrho_n.$$

Since Λ is a v -perturbation, we therefore get

$$\sum_{n=1}^{\infty} |a_n|^2 v_n \lesssim \sum_{n=1}^{\infty} \frac{\varrho_n}{|\gamma_n|^2 V_n} < \infty,$$

where in the final step we used that the ratio ϱ_n/V_n grows at most sub-exponentially. We then get

$$|h(z)|^2 \lesssim \frac{\varrho_n}{|z|^4} + \frac{V_n}{|z|^2}$$

when z is in $D_n(v; M)$ with M sufficiently large. Using again that both ϱ_n and V_n grow at most sub-exponentially, we have that $h(z) \rightarrow 0$ when $z \rightarrow \infty$, which means that $h \equiv 0$. \square

LEMMA 5.8. *Suppose that $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$. If, in addition, Λ is a v -perturbation of Γ of deficiency 1, then Λ is not a uniqueness sequence for $H_{(\Gamma,v)}$.*

Proof. In this case, we may write

$$\frac{c}{z - \gamma_1} \prod_{m=2}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_m} = \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} + h(z),$$

where h is an entire function and

$$|a_n|^2 v_n^2 \simeq \frac{|\gamma_n - \lambda_n|^2}{|\gamma_n|^2} \varrho_n.$$

Since Λ is a v -perturbation, we get

$$\sum_{n=1}^{\infty} |a_n|^2 v_n \lesssim \sum_{n=1}^{\infty} \frac{\varrho_n}{|\gamma_n|^2 P_n} < \infty,$$

where we now used that the ratio ϱ_n/P_n grows at most sub-exponentially. It follows that

$$|h(z)|^2 \lesssim \frac{\varrho_n}{|z|^2} + P_n$$

when z is in $D_n(v; M)$ with M sufficiently large. We conclude that $h(z) \rightarrow 0$ when $z \rightarrow \infty$, which means that $h \equiv 0$. \square

5.3. Geometric criteria for invertibility of $H_{(\Gamma, v); (\Lambda, w)}$ when Γ is a sparse sequence

After the preliminary results of the previous subsection, we may now state our geometric conditions for invertibility. We begin with the case when $v_n = o(V_n)$ as $n \rightarrow \infty$.

THEOREM 5.1. *Suppose that w is the Bessel weight sequence for Λ with respect to (Γ, v) and that $V_n \rightarrow \infty$ and $v_n = o(V_n)$ when $n \rightarrow \infty$. Then the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is invertible if and only if $\sup_n V_n Q_n < \infty$ and one of the following conditions holds.*

(1) *The sequence Λ is an exact v -perturbation of Γ and there are positive constants C and δ such that*

$$\frac{\varrho_m}{\varrho_n} \leq C \left(\frac{V_m}{V_n} \right)^{1-\delta} \quad (5.5)$$

whenever $m > n$.

(2) *The sequence Λ is a v -perturbation of Γ of defect 1 and there are positive constants C and δ such that*

$$\frac{\varrho_m}{\varrho_n} \geq C \left(\frac{V_m}{V_n} \right)^{1+\delta} \quad (5.6)$$

whenever $m > n$.

It is quite remarkable that the essential quantitative conditions for invertibility, found in (1) and (2), only depend on the moduli of the complex numbers γ_n/λ_n .

We note that in the case when

$$\sum_{n=1}^{\infty} v_n < \infty,$$

the result is much simpler and less delicate. Then, as can be seen from the proof of part (1) of Theorem 5.1, the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is invertible if and only if Λ is an exact v -perturbation of Γ and $\sup_n Q_n < \infty$. This result can be viewed as a special case of part (1) of the theorem.

To arrive at the results stated in Example 1 (see Section 1), we note that if

$$\frac{|\gamma_n - \lambda_n|}{|\lambda_n|} \lesssim \frac{v_n}{V_n},$$

then

$$Q_n = \sum_{m=n+1}^{\infty} \frac{w_m}{|\lambda_m|^2} \lesssim \sum_{m=n+1}^{\infty} \frac{v_m}{V_{m+1}^2} \leq \frac{1}{V_{n+1}},$$

where in the last step we compared the sum with the integral of $1/x^2$ from V_{n+1} to ∞ . We also have, assuming $|\gamma_n|/|\lambda_n| - 1 \leq cv_n/V_n$, that

$$\log \frac{\varrho_m}{\varrho_n} \leq 2c(1 + o(1)) \sum_{j=n+1}^m \frac{v_l}{V_l} = 2c(1 + o(1)) \log \frac{V_m}{V_n} \quad (5.7)$$

when $m > n$ and $n \rightarrow \infty$. In view of Theorem 5.1, this gives part (1) of the example; part (2) follows by the same argument, with the inequality in (5.7) reversed.

In the case when $v_n/|\gamma_n|^2 = o(P_n)$, we have the following counterpart to Theorem 5.1.

THEOREM 5.2. *Suppose that w is the Bessel weight sequence for Λ with respect to (Γ, v) and that $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$. Then the transformation $H_{(\Gamma, v); (\Lambda, w)}$ is invertible if and only if we have $\sup_n W_n P_n < \infty$ and one of the following two conditions holds.*

(1) *The sequence Λ is an exact v -perturbation of Γ and there are positive constants C and δ such that*

$$\frac{\varrho_m}{\varrho_n} \geq C \left(\frac{P_m}{P_n} \right)^{1-\delta} \quad (5.8)$$

whenever $m > n$.

(2) *The sequence Λ is a v -perturbation of Γ of excess 1 and there are positive constants C and δ such that*

$$\frac{\varrho_m}{\varrho_n} \leq C \left(\frac{P_m}{P_n} \right)^{1+\delta} \quad (5.9)$$

whenever $m > n$.

There is a slight lack of symmetry between the two theorems; while it may happen that $\sup_n V_n < \infty$, we will always have that $P_n \rightarrow 0$. Therefore, no precaution is needed concerning the decay of P_n .

We have the following statement, in complete analogy with Example 1 and with the same proof.

EXAMPLE 2. Suppose that $v_n/|\gamma_n|^2 = o(P_n)$ when $n \rightarrow \infty$ and that $\sup_n W_n P_n < \infty$.

(1) If, in addition, Λ is an exact v -perturbation of Γ and there is a real constant $c < 1/2$ such that

$$\frac{|\lambda_n|}{|\gamma_n|} - 1 \leq c \frac{v_n}{|\gamma_n|^2 P_n}$$

for all sufficiently large n , then $H_{(\Gamma, v); (\Lambda, w)}$ is an invertible transformation.

(2) If, on the other hand, Λ is a v -perturbation of Γ of excess 1 and there is a positive constant $c > 1/2$ such that

$$\frac{|\lambda_n|}{|\gamma_n|} - 1 \geq c \frac{v_n}{|\gamma_n|^2 P_n}$$

for all sufficiently large n , then $H_{(\Gamma, v); (\Lambda, w)}$ is an invertible transformation.

The two final subsections of this paper will present the proof of Theorem 5.1; the proof of Theorem 5.2 is completely analogous and will therefore be omitted.

5.4. Proof of the necessity of the conditions in Theorem 5.1

In addition to the results of Subsection 5.2, we need the following simple facts.

LEMMA 5.9. *Let $c = (c_n)$ be a sequence of positive numbers.*

(i) *If there is a constant C such that $\sum_{m=1}^{n-1} c_m \leq Cc_n$ for $n > 1$, then there is a positive constant δ such that $c_m/c_n \geq C2^{\delta(m-n)}$ whenever $m > n$.*

(ii) *If there is a constant C such that $\sum_{m=n+1}^{\infty} c_m \leq Cc_n$ for every positive integer n , then there is a positive constant δ such that $c_m/c_n \leq C2^{-\delta(m-n)}$ whenever $m > n$.*

Proof. We consider (i). The assumption implies that

$$Nc_{n-1} \leq N \sum_{m=1}^{n-1} c_m \leq C \sum_{m=n}^{n+N-1} c_m \leq C^2 c_{n+N}.$$

which means that if we choose $N > 2C^2$, then $c_{n+j(N+1)} \geq 2^j c_n$. The result follows if we choose $\delta = 1/(N+2)$. The proof of (ii) can be performed in a similar way. \square

We turn to the proof of the necessity of the conditions in Theorem 5.1. Thus we begin by assuming that $H_{(\Gamma,v);(\Lambda,w)}$ is an invertible transformation. Since this means that, in particular, $H_{(\Gamma,v);(\Lambda,w)}$ is a bounded transformation, we must have $\sup_n V_n Q_n < \infty$. Also, in view of Lemma 5.1, we already know that Λ is either an exact v -perturbation of Γ or a v -perturbation of Γ of deficiency 1. Thus it remains only to establish the necessity of the conditions in parts (1) and (2), under the respective assumptions of exactness and deficiency 1.

We treat the two cases separately.

Case 1: Λ is an exact v -perturbation of Γ . Since $v_n = o(V_n)$, the weight sequence $w = (w_n)$ defined by (1.7) satisfies

$$w_n \simeq \frac{|\gamma_n - \lambda_n|^2}{v_n}. \quad (5.10)$$

As a consequence, we now obtain simple estimates for the weight sequences $\nu = (\nu_n)$ and $\varpi = (\varpi_j)$ appearing in Theorem 1.3.

We begin by noting that if Λ is a v -perturbation of Γ and an exact uniqueness sequence for $H_{(\Gamma,v)}$, then there is a constant c such that

$$\sum_{n=1}^{\infty} \frac{e_n v_n}{z - \gamma_n} = \frac{c}{z - \gamma_1} \prod_{m=2}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_m} \quad (5.11)$$

for every z in $\mathbb{C} \setminus \Gamma$, where again $e = (e_n)$ is the vector such that $H_{(\Gamma,v);(\Lambda,w)} e = (1, 0, 0, \dots)$. Indeed, the expression on the left-hand side can have zeros only at the points λ_m for $m > 1$, since Λ is assumed to be an exact uniqueness sequence for $H_{(\Gamma,v)}$. From (5.11) we obtain

$$|e_n|^2 v_n^2 \simeq \frac{|\lambda_n - \gamma_n|^2}{|\lambda_n|^2} \varrho_n,$$

and therefore, using (1.8) and (5.10), we obtain

$$\nu_n \simeq w_n \varrho_n. \quad (5.12)$$

On the other hand, differentiating (5.11) at $z = \lambda_n$, we get

$$\left| \sum_{l=1}^{\infty} \frac{e_l v_l}{(\lambda_n - \gamma_l)^2} \right| \simeq \frac{|\gamma_n|}{|\lambda_n|^2 |\lambda_n - \gamma_n|} \prod_{m=1}^{n-1} \frac{|\gamma_m|}{|\lambda_m|}.$$

Thus using (1.9) and again (5.10), we obtain

$$\varpi_n \simeq v_n \varrho_n^{-1}. \quad (5.13)$$

To simplify the writing, we set

$$V_n^{(\varrho,0)} = \sum_{m=1}^{n-1} v_m \varrho_m^{-1} \quad \text{and} \quad P_n^{(\varrho,0)} = \sum_{m=n+1}^{\infty} v_m |\lambda_m|^{-2} \varrho_m^{-1}$$

as well as

$$W_n^{(\varrho,0)} = \sum_{m=1}^{n-1} w_m \varrho_m \quad \text{and} \quad Q_n^{(\varrho,0)} = \sum_{m=n+1}^{\infty} w_m \varrho_m |\lambda_n|^{-2}.$$

By Theorems 1.1 and 1.3, we must have $\sup_n V_n^{(\varrho,0)} Q_n^{(\varrho,0)} < \infty$; we will now show that the estimate in part (1) is a consequence of this condition.

We set $n_1 = 2$ and define n_j inductively by requiring $V_{n_{j+1}-1}/V_{n_j} < 2 \leq V_{n_{j+1}}/V_{n_j}$. By (5.2) of Lemma 5.3 and the uniform boundedness of $V_n Q_n$, it follows that there are constants c and C such that $c < \varrho_n/\varrho_m \leq C$ when n and m both lie in the interval $[n_j, n_{j+1}]$. Hence we have

$$V_{n_j}^{(\varrho,0)} \simeq \sum_{l=1}^j V_{n_l} \varrho_{n_l}^{-1}. \quad (5.14)$$

Now if

$$Q_{n_j} - Q_{n_{j+1}} \geq \frac{\varepsilon}{V_{n_{j+1}}}, \quad (5.15)$$

then our condition $\sup_n V_n^{(\varrho,0)} Q_n^{(\varrho,0)} < \infty$ and (5.14) imply that there exists a constant C such that

$$\sum_{l=1}^j V_{n_l} \varrho_{n_l}^{-1} \leq C V_{n_{j+1}} \varrho_{n_{j+1}}^{-1}. \quad (5.16)$$

If, on the other hand, we have

$$Q_{n_j} - Q_{n_{j+1}} < \frac{\varepsilon}{V_{n_{j+1}}},$$

then an application of (5.1) of Lemma 5.3 gives $\varrho_{n_{j+1}}/\varrho_{n_j} \leq 5/4$ if ε is sufficiently small. Hence we have

$$\frac{V_{n_{j+1}} \varrho_{n_j}}{V_{n_j} \varrho_{n_{j+1}}} \geq \frac{8}{5},$$

which means that $V_{n_j} \varrho_{n_j}^{-1}$ increases exponentially on any set of consecutive integers j for which (5.15) fails. Combining (5.16) with the latter estimate, we therefore get that

$$\sum_{l=1}^j V_{n_l} \varrho_{n_l}^{-1} \leq \left(\frac{5}{8}C + \frac{8}{3} \right) V_{n_{j+1}} \varrho_{n_{j+1}}^{-1}$$

when (5.15) fails and ε is sufficiently small. Thus (5.16) holds for every index j if the constant C is suitably adjusted. Hence, by part (i) of Lemma 5.9, there exists a constant C such that

$$\frac{\varrho_{n_{j+l}}}{\varrho_{n_j}} \leq C \frac{V_{n_{j+l}}}{V_{n_j}} 2^{-\delta l} \leq C \left(\frac{V_{n_{j+l}}}{V_{n_j}} \right)^{1-\delta/2},$$

where in the last step we used that $V_{n_{j+1}}/V_{n_j} \leq 4$ for sufficiently large j . We are done since it suffices to establish (5.5) for $n = n_j$ and $m = n_{j+l}$.

Case 2: Λ is a v -perturbation of Γ of deficiency 1. As in the previous case, we begin by finding estimates for the weight sequences $\nu = (\nu_n)$ and $\varpi = (\varpi_j)$ appearing in Theorem 1.3. If Λ is a v -perturbation of Γ of deficiency 1 and an exact uniqueness sequence for $H_{(\Gamma,v)}$, then

there is a constant c such that

$$\sum_{n=1}^{\infty} \frac{e_n v_n}{z - \gamma_n} = \frac{c}{(z - \gamma_1)(z - \gamma_2)} \prod_{m=3}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_n}$$

for every z in $\mathbb{C} \setminus \Gamma$, where again $e = (e_n)$ is the vector such that $H_{(\Gamma, v); (\Lambda, w)} e = (1, 0, 0, \dots)$. Arguing in the same way as in the preceding case, we obtain from this relation the estimates

$$\nu_n \simeq w_n \varrho_n |\gamma_n|^{-2} \quad (5.17)$$

and

$$\varpi_n \simeq v_n \varrho_n^{-1} |\gamma_n|^2. \quad (5.18)$$

We now set

$$V_n^{(\varrho, 1)} = \sum_{m=1}^{n-1} v_n |\gamma_n|^2 \varrho_n^{-1} \quad \text{and} \quad P_n^{(\varrho, 1)} = \sum_{m=n+1}^{\infty} v_n \varrho_n$$

as well as

$$W_n^{(\varrho, 1)} = \sum_{m=1}^{n-1} w_n |\gamma_n|^{-2} \varrho_n \quad \text{and} \quad Q_n^{(\varrho, 1)} = \sum_{m=n+1}^{\infty} w_n |\gamma_n|^{-4} \varrho_n.$$

By Theorems 1.1 and 1.3, we must have $\sup_n W_n^{(\varrho, 1)} P_n^{(\varrho, 1)} < \infty$; we will now show that also the estimate in part (2) is a consequence of this condition.

We let the sequence $(n_j)_j$ be as above and find that

$$P_{n_j}^{(\varrho, 1)} \simeq \sum_{l=j+1}^{\infty} V_{n_l} \varrho_{n_l}^{-1} \quad (5.19)$$

whenever $j \geq 1$. Now if

$$Q_{n_{j+1}} - Q_{n_j} \geq \frac{\varepsilon}{V_{n_{j+1}}}, \quad (5.20)$$

then it follows from the condition $\sup_n W_n^{(\varrho, 1)} P_n^{(\varrho, 1)} < \infty$ and (5.19) that

$$\sum_{l=j+1}^{\infty} V_{n_l} \varrho_{n_l}^{-1} \lesssim V_{n_j} \varrho_{n_j}^{-1}. \quad (5.21)$$

As in the preceding case, we find that if ε is sufficiently small, then $V_{n_j} \varrho_{n_j}^{-1}$ increases exponentially on any set of consecutive integers j for which (5.20) fails. The relation (5.19) implies that no such set is infinite; thus there is an infinite sequence of indices n_j for which (5.21) holds, and there must in fact be a uniform bound on the number of points found in any set of consecutive integers j for which (5.20) fails. We may infer from this argument that in fact (5.21) holds for every index $n_j \geq 1$. Finally, we invoke part (ii) of Lemma 5.9, which implies that there is a constant C such that

$$\frac{\varrho_{n_{j+l}}}{\varrho_{n_j}} \geq C \frac{V_{n_{j+l}}}{V_{n_j}} 2^{\delta l} \geq C \left(\frac{V_{n_{j+l}}}{V_{n_j}} \right)^{1+\delta},$$

and we are done since it suffices to establish (5.6) for $n = n_j$ and $m = n_{j+l}$.

5.5. Proof of the sufficiency of the conditions in Theorem 5.1

We begin by noting that the condition $\sup_n V_n Q_n < \infty$ implies that $H_{(\Gamma, v); (\Lambda, w)}$ is a bounded transformation. Indeed, (1.4) in Theorem 1.1 holds trivially when $\mu = \sum_n w_n \delta_{\lambda_n}$. We also have

$$W_n \lesssim \frac{|\gamma_n|^2}{V_n} \quad \text{and} \quad P_n \lesssim \frac{v_n}{|\gamma_n|^2}$$

by the assumptions that $v_n = o(V_n)$ and $\sup_n V_n Q_n < \infty$. Therefore, Theorem 1.1 allows us to conclude that $H_{(\Gamma, v); (\Lambda, w)}$ is a bounded transformation.

We now use Theorem 1.3 and show that the respective conditions in parts (1) and (2) in Theorem 5.1 imply those in Theorem 1.3. The sequence $(n_j)_j$ will be the same as in the previous subsection.

Case 1: Λ is an exact v -perturbation of Γ . We already know from Lemma 5.5 that if Λ is an exact v -perturbation of Γ , then Λ is a uniqueness sequence for $H_{(\Gamma, v)}$. To check that Λ is in fact an exact uniqueness sequence for $H_{(\Gamma, v)}$, we note that we may write

$$\frac{c}{z - \gamma_1} \prod_{m=2}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_n} = \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} + h(z),$$

where h is an entire function and

$$|a_n|^2 v_n \simeq \frac{w_n}{|\gamma_n|^2} \varrho_n.$$

By the assumption that $\sup_n V_n Q_n < \infty$, we have

$$\sum_{n=1}^{\infty} |a_n|^2 v_n \lesssim \sum_{j=1}^{\infty} \frac{\varrho_{n_j}}{V_{n_j}},$$

which, in view of (5.5), implies that (a_n) is in ℓ_v^2 . In particular, we then have

$$|h(z)|^2 \lesssim \frac{\varrho_n}{|z|^2} + \frac{V_n}{|z|^2}$$

when z is in $D_n(v; M)$ with M sufficiently large. Thus $h(z) \rightarrow 0$ when $z \rightarrow \infty$, which means that $h \equiv 0$.

It remains only to verify that $H_{(\Lambda, \varpi); (\Gamma, \nu)}$ is a bounded transformation. By Theorem 1.1, we need to show that we have both $\sup_n W_n^{(e,0)} P_n^{(e,0)} < \infty$ and $\sup_n V_n^{(e,0)} Q_n^{(e,0)} < \infty$. To this end, we note that since ϱ_n can only grow sub-exponentially, we have $\sup_n W_n^{(e,0)} P_n^{(e,0)} < \infty$ by the same argument that gave $\sup_n W_n P_n < \infty$. Since $\sup_n V_n Q_n < \infty$, we have

$$V_n^{(e,0)} Q_n^{(e,0)} \lesssim \sum_{n_j < n} \frac{V_{n_j}}{\varrho_{n_j}} \frac{\varrho_n}{V_n};$$

here the right-hand side is uniformly bounded whenever (5.5) holds.

Case 2: Λ is a v -perturbation of Γ of deficiency 1. In view of Lemma 5.7, we have that Λ is an exact uniqueness sequence for $H_{(\Gamma, v)}$ if we can show that there is no nonzero a in ℓ_v^2 such that $H_{(\Gamma, v)} a$ vanishes on Λ . To show this, we assume to the contrary that such a sequence a exists. Then there is a constant c such that

$$\sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} = \frac{c}{z - \gamma_1} \prod_{m=2}^{\infty} \frac{1 - z/\lambda_m}{1 - z/\gamma_m}. \quad (5.22)$$

By estimating each side of (5.22) for z in $D_n(v; M)$ with M sufficiently large, we get

$$V_n \sum_{m=1}^{\infty} |a_m|^2 v_m \gtrsim \varrho_n.$$

But this is a contradiction, because (5.6) implies that ϱ_n/V_n is an increasing sequence.

It remains only to verify that $H_{(\Lambda, \varpi); (\Gamma, \nu)}$ is a bounded transformation. To this end, we note that $\sup_n V_n^{(e,1)} Q_n^{(e,1)} < \infty$ holds trivially because $1/\varrho_n$ can only grow sub-exponentially, while

$$W_n^{(e,1)} P_n^{(e,1)} \lesssim \sum_{n_j < n} \frac{P_{n_j}}{\varrho_{n_j}} \frac{\varrho_n}{P_n},$$

which is uniformly bounded when (5.6) holds.

References

1. P. R. AHERN and D. N. CLARK, 'Radial limits and invariant subspaces', *Amer. J. Math.* 92 (1970) 332–342.
2. A. B. ALEKSANDROV, 'A simple proof of the Volberg–Treil theorem on the embedding of coinvariant subspaces of the shift operator', *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 217 (1994), *Issled. po Linein. Oper. i Teor. Funktsii.* 22, 26–35, 218 (Russian), *J. Math. Sci. (N. Y.)* 85 (1997) 1773–1778 (English).
3. A. B. ALEKSANDROV, 'Embedding theorems for coinvariant subspaces of the shift operator II', *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 262 (1999), *Issled. po Linein. Oper. i Teor. Funktsii.* 27, 5–48, 231 (Russian), *J. Math. Sci. (N. Y.)* 110 (2002) 2907–2929.
4. A. D. BARANOV, 'Bernstein-type inequalities for shift-coinvariant subspaces and their applications to Carleson embeddings', *J. Funct. Anal.* 223 (2005) 116–146.
5. A. D. BARANOV, 'Stability of bases and frames of reproducing kernels in model subspaces', *Ann. Inst. Fourier (Grenoble)* 55 (2005) 2399–2422.
6. A. BARANOV and K. DYAKONOV, 'The Feichtinger conjecture for reproducing kernels in model subspaces', *J. Geom. Anal.* (2010), DOI: 10.1007/s12220-010-9147-y.
7. Y. BELOV, T. MENGESTIE and K. SEIP, 'Unitary discrete Hilbert transforms', *J. Anal. Math.* 112 (2010) 383–393.
8. A. BEURLING, 'On two problems concerning linear transformations in Hilbert space', *Acta Math.* 81 (1948) 239–255.
9. A. BORICHEV and YU. LYUBARSKII, 'Riesz bases of reproducing kernels in Fock-type spaces', *J. Inst. Math. Jussieu* 9 (2010) 449–461.
10. L. DE BRANGES, *Hilbert spaces of entire functions* (Prentice-Hall, Englewood Cliffs, 1968).
11. P. G. CASAZZA, O. CHRISTENSEN, A. LINDNER and R. VERSHYNIN, 'Frames and the Feichtinger conjecture', *Proc. Amer. Math. Soc.* 133 (2005) 1025–1033.
12. P. G. CASAZZA, M. FICKUS, J. C. TREMAIN and E. WEBER, 'The Kadison–Singer problem in mathematics and engineering: a detailed account', *Operator theory, operator algebras, and applications*, Contemporary Mathematics 414 (American Mathematical Society, Providence, RI, 2006) 299–356.
13. P. G. CASAZZA, G. KUTINYOK, D. SPEEGLE and J. C. TREMAIN, 'A decomposition theorem for frames and the Feichtinger conjecture', *Proc. Amer. Math. Soc.* 136 (2008) 2043–2053.
14. P. G. CASAZZA and J. C. TREMAIN, 'The Kadison–Singer problem in mathematics and engineering', *Proc. Natl. Acad. Sci. USA* 103 (2006) 2032–2039.
15. P. G. CASAZZA and R. VERSHYNIN, 'Kadison–Singer meets Bourgain–Tzafriri', Preprint, 2005.
16. D. N. CLARK, 'One dimensional perturbations of restricted shifts', *J. Anal. Math.* 25 (1972) 169–191.
17. W. COHN, 'Carleson measures for functions orthogonal to invariant subspaces', *Pacific J. Math.* 103 (1982) 347–364.
18. W. COHN, 'Radial embedding theorems for invariant subspaces', *Complex Var. Theory Appl.* 17 (1991) 33–42.
19. K. M. DYAKONOV, 'Embedding theorems for star-invariant subspaces generated by smooth inner functions', *J. Funct. Anal.* 157 (1998) 588–598.
20. S. V. HRUŠČEV, N. K. NIKOL'SKIĬ and B. S. PAVLOV, 'Unconditional bases of exponentials and of reproducing kernels', *Complex analysis and spectral theory* (eds V. P. Havin and N. K. Nikol'skiĭ), Lecture Notes in Mathematics 864 (Springer, Berlin, 1981) 214–335.
21. M. I. KADETS, 'The exact value of the Paley–Wiener constant', *Sov. Math. Dokl.* 5 (1964) 559–561.
22. R. KADISON and I. SINGER, 'Extensions of pure states', *Amer. J. Math.* 81 (1959) 383–400.
23. YU. LYUBARSKII and K. SEIP, 'Complete interpolating sequences and Muckenhoupt's (A_p) condition', *Rev. Mat. Iberoamericana* 13 (1997) 361–376.
24. YU. LYUBARSKII and K. SEIP, 'Weighted Paley–Wiener spaces', *J. Amer. Math. Soc.* 15 (2002) 979–1006.
25. F. NAZAROV and A. VOLBERG, 'Bellman function, two weighted Hilbert transform, and embeddings of the model space K_Θ ', *J. Anal. Math.* 87 (2002) 385–414.
26. N. K. NIKOL'SKIĬ, *Treatise on the shift operator*, Grundlehren der Mathematischen Wissenschaften 273 (Springer, Berlin, 1986).
27. N. K. NIKOLSKI, *Operators, functions, and systems: an easy reading*, vol. 1, Mathematical Surveys and Monographs 92 (American Mathematical Society, Providence, RI, 2002).
28. N. K. NIKOLSKI, *Operators, functions, and systems: an easy reading*, vol. 2, Mathematical Surveys and Monographs 92 (American Mathematical Society, Providence, RI, 2002).
29. W. RUDIN, *Functional analysis* (McGraw-Hill, New York, 1973).
30. S. R. TREIL and A. L. VOLBERG, 'Embedding theorems for invariant subspaces of the inverse shift operator', *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 149 (1986) 38–51 (Russian), *J. Soviet Math.* 42 (1988) 1562–1572 (English).
31. A. L. VOLBERG, *Calderón–Zygmund capacities and operators on nonhomogeneous spaces*, CBMS Regional Conference Series in Mathematics 100 (American Mathematical Society, Providence, RI, 2003).

Yurii Belov, Tesfa Y. Mengestie and Kristian Seip
Department of Mathematical Sciences
Norwegian University of Science and Technology (NTNU)
NO-7491 Trondheim
Norway

j_b_juri_belov@mail.ru
mengesti@math.ntnu.no
seip@math.ntnu.no